arXiv:1110.6794v1 [gr-qc] 31 Oct 2011

Decay properties of Klein-Gordon fields on Kerr-AdS spacetimes

Gustav Holzegel* and Jacques Smulevici†

November 1, 2011

Abstract

This paper investigates the decay properties of solutions to the massive linear wave equation $\Box_g \psi + \frac{\alpha}{l^2}\psi = 0$ for g the metric of a Kerr-AdS spacetime satisfying $|a|l < r_+^2$ and $\alpha < \frac{9}{4}$ satisfying the Breitenlohner Freedman bound. We prove that the non-degenerate energy of ψ with respect to an appropriate foliation of spacelike slices decays like $(\log t^\star)^{-1}$. Our estimates are expected to be sharp from heuristic and numerical arguments in the physics literature suggesting that general solutions will only decay logarithmically. The underlying reason for the slow decay rate can be traced back to a stable trapping phenomenon for asymptotically anti de Sitter black holes which is in turn a consequence of the reflecting boundary conditions for ψ at null-infinity.

Contents

1 Introduction 3
1.1 Logarithmic Decay 4
1.2 Absence of time-periodic solutions 5
1.3 Quasi-normal modes 6
1.4 The non-linear stability problem 6
1.5 Outline 7

2 The Kerr-AdS family of spacetimes 8
2.1 The fixed manifold with boundary $\mathcal{R}$ 8
2.2 The parameter space 8
2.3 Kerr-AdS-star coordinates 8
2.4 The tortoise coordinate r^* 9
2.5 Boyer-Lindquist coordinates 9
2.6 The Kerr-AdS metric for fixed (a, M, l) 9

---

[*] Princeton University, Department of Mathematics, Fine Hall, Washington Road, Princeton, NJ 08544, United States.

[†] Laboratoire de Mathématiques, Université Paris-Sud 11, bât. 425, 91405 Orsay, France.

2.7 Coordinate Vectorfields and Isometries 11
2.8 Foliations 11

3 The norms 11

4 Well-posedness and boundedness statement 12

5 Preliminaries 14
5.1 Time cut-off 14
5.2 The (modified) oblate spheroidal harmonics 15
5.3 The separation of variables 17
5.4 Properties of the Potential 18
5.5 (ℓ, m, ω) low-frequency localisation 21

6 Microlocal current templates 22
6.1 Microlocal energy and red-shift currents 22
6.2 Microlocal Morawetz currents 22

7 Microlocal estimates for $\psi^\tau_{\flat,\omega\approx\omega_+}$ 23

8 Microlocal estimates for $\psi^\tau_{\flat,\omega\not\approx\omega_+}$ 26
8.1 Preliminaries 27
  8.1.1 Exponentially weighted multipliers 27
  8.1.2 An exponentially weighted Hardy inequality 28
8.2 Estimating $|u'|^2$ from $|u|^2$ 29
8.3 Estimating $|u|^2$ from $|u'|^2$ 32
8.4 Putting it together: The basic estimate 32

9 Spacetime estimate for $\psi^\tau_\flat = \psi^\tau_{\flat,\omega\approx\omega_+} + \psi^\tau_{\flat,\omega\not\approx\omega_+}$ 33
9.1 The main terms 34
9.2 The H-error-terms 34
9.3 The boundary-terms 35
9.4 Improving the weights near null-infinity 36
9.5 Improving the weights near the horizon 37
9.6 Higher derivatives 38

10 Estimates for $\psi^\tau_\sharp$ 39

11 Proof of Theorem 1.1: Logarithmic decay 39

12 Some remarks on the Schwarzschild-AdS case 41

13 Proof of Theorem 1.3: Absence of periodic solutions 42

14 Acknowledgments 45

A Basic Fourier estimates 45

1 Introduction

In a sequence of recent papers [27, 26, 28, 29], the study of boundedness and decay properties of waves on asymptotically Anti-de Sitter (AdS) black hole spacetimes was initiated. The analysis focused on the dynamics of the massive wave equation

$$\Box_g \psi + \frac{\alpha}{l^2}\psi = 0 \tag{1}$$

for g an asymptotically AdS spacetime with cosmological constant $\Lambda = -3l^{-2}$ and α satisfying $\alpha < \frac{9}{4}$, the Breitenlohner-Freedman bound. The well-posedness of this equation on such backgrounds was established in [34, 27] under suitable boundary conditions at null-infinity. (See also [4] for earlier results on exactly AdS.) The uniform boundedness of solutions to (1) was proven in [26] for Kerr-AdS spacetimes, provided the angular momentum per unit mass of the black hole is not too large.[1] The proof relied on three crucial ingredients:

1. the existence of a globally causal vectorfield on the black hole exterior (the Hawking-Reall vectorfield [24]) for the range $|a|l < r_+$ where $r_+ = r_+(M, a, l)$ is the r-value at the horizon in Boyer-Lindquist coordinates,
2. the exploitation of weighted Hardy inequalities allowing one to absorb the (wrong-signed) mass-term in the range $0 < \alpha < \frac{9}{4}$,
3. the redshift effect near the event horizon coupled with the Hawking-Reall vectorfield allowed to prove boundedness *without* the construction of an integrated decay (Morawetz) estimate. This goes back to an idea of Dafermos and Rodnianski in the asymptotically-flat, Schwarzschild-case, cf. [15].

The ingredients 1. and 3. are particularly remarkable, when compared with the situation in the asymptotically flat case, where the boundedness statement for solutions of the massless wave equation on Kerr with $a \ll M$ is already intimately connected with the decay properties of the solution [20]: The absence of a globally causal Killing field implies that techniques based purely on spacetime vectorfields are insufficient to control the solution and instead microlocal analysis is required (or higher order differential operators, see [2])). Interestingly, integrated *decay* estimates had to be constructed for superradiant frequency ranges to prove *boundedness* of the solution. The Kerr-AdS-case, on the other hand, is (precisely because of (1) and (3)) more similar to the asymptotically-flat Schwarzschild case, where the boundedness and the decay statements can also be entirely decoupled.

Regarding 1. above, we also remark that in the range $|a|l > r_+$ solutions are expected to grow because superradiant effects are continuously amplified by the reflecting AdS boundary [10, 12, 11].

---

[1] In fact, the result was shown to hold for a class of C^1 perturbations of these spacetimes.

While the decay of solutions on asymptotically-flat Kerr black holes has recently been understood for the subextremal range $|a| < M$ [19] after a period of intense research over the last ten years [14, 17, 15, 19, 18, 16, 31, 33, 2], the question of decay of solutions to (1) remained unanswered in [26]. Very recently the authors, in the context of a spherically symmetric non-linear problem [28, 29], proved in particular that spherically symmetric solutions of (1) decay exponentially on the Schwarzschild-AdS black hole exterior provided $\alpha \leq 2$ holds. The latter result exploited a version of the redshift near null-infinity, as well as the absence of any trapping phenomena in spherical symmetry.

Beyond spherically symmetric solutions of the wave equation, however, the picture changes radically. This may be anticipated at the heuristic level from an analysis of geodesic flow (which is the high frequency or "geometric optics" approximation for the propagation of waves) on the Schwarzschild-AdS manifold (i.e. $a = 0$). As in the asymptotically flat case, the radial equation for the null-geodesics sees a potential $V(r)$ which exhibits a local maximum at $r = 3M$ corresponding to the familiar photon sphere. Consequently, ingoing geodesics which do not have sufficient energy to surmount the potential will be reflected off towards null-infinity. While in the asymptotically flat case these (now outgoing) geodesics eventually leave the spacetime through null-infinity, in the asymptotically AdS case, they are reflected back into the spacetime. This is a stable trapping phenomenon, which constitutes a serious obstacle to prove decay. The above reasoning does not take ' 'tunneling-effects" (i.e. the uncertainty principle for the wave equation) into account. Indeed, an exponentially small part of the energy will tunnel through the potential and leave the black hole exterior through the event horizon. From these heuristics one may expect a logarithmic decay rate and this is indeed what we shall prove below.

1.1 Logarithmic Decay

We now state our main results directing the reader to Section 3 for a precise definition of the energy norms and functional spaces involved in Theorem 1.1. The statement below also makes reference to a foliation of the Kerr-AdS black hole exterior by spacelike slices, $\Sigma_{t^\star}$, which will be defined in Section 2. For the reader familiar with the formalism of Penrose diagrams, the foliation may be read off from Figure 1.

Theorem 1.1. *Let* $(\mathcal{M}, g)$ *be a Kerr-AdS spacetime with parameters* (M, a, l) *for which the boundedness theorem of [26] holds*[2] *and* ψ *be a* CH^2_{AdS} *solution of (1) with* $-\infty < \alpha < \frac{9}{4}$ *on this background. Assume moreover that one of the following three conditions on the parameters hold:*

1. $\alpha \leq 1$,
2. $1 < \alpha \leq 2$ *and* $|a| < \frac{l}{2}$,

---

[2] In particular, both the Hawking-Reall condition $r_+^2 > |a|l$ and the no-naked singularities condition $|a| < l$ hold.

3. a is sufficiently small depending only on M, l and α.

Then, for the foliation of spacelike slices $\Sigma_{t^\star}$ (intersecting the event horizon) defined in Section 2.8, we have the decay estimate

$$\|\psi\|^2_{H^1_{AdS}(\Sigma_{t^\star})} \lesssim (\log\ t^\star)^{-1} \left[\|\psi\|^2_{H^2_{AdS}(\Sigma_0)}\right] \tag{2}$$

for any $t^\star \geq 2$, where $\lesssim$ allows a constant depending only on the parameters M, a, l of the background spacetime and on α.

By what have become standard techniques (commuting with T, the redshift and angular momentum vectorfields, cf. [15], [26]), one obtains this $\log\ t^\star$-decay for all higher order norms and also pointwise estimates via Sobolev embedding. In particular, one has

Corollary 1.2. *Under the assumptions of Theorem 1 and with ψ in addition in CH^3_{AdS} one has the pointwise bound*

$$|\psi|^2 \lesssim (\log\ t^\star)^{-1} \left[\|\psi\|^2_{H^3_{AdS}(\Sigma_0)}\right] . \tag{3}$$

We remark that we have made no serious attempt to optimize the conditions on the parameter a in items 2 and 3 of Theorem 1.1. They can most likely be improved with further work.



Figure 1: The spacelike hypersurfaces Σ_t and $\Sigma_{t^\star}$.

1.2 Absence of time-periodic solutions

Obviously, Theorem 1.1 excludes time-periodic solutions. This situation is in sharp contrast with that of pure AdS, where periodic solutions may be easily constructed, cf. [8]. It turns out that the absence of periodic solutions can be proven in a more elementary fashion which is why we include and prove this as a separate theorem.

Theorem 1.3. *Let* $(\mathcal{M}, g)$ *be a Kerr-AdS spacetime with parameters* (M, a, l) *such that* $|a|l < r_+^2$ *and* $|a| < l$*. Let* ψ *be a solution of (1) in the class* CH^1_{AdS}*, with* $\alpha < 9/4$*, with* ψ *of the form* $\psi = e^{i\omega t}\hat{\psi}$*, where* $\omega \in \mathbb{R}$ *and* $\hat{\psi}$ *is constant in* t*. Assume moreover that one of the three conditions on the parameters given in Theorem 1.1 hold. Then* $\psi = 0$*.*

The above theorem implies the absence of time-periodic solutions in CH^1_{AdS}: Indeed, given a solution ψ in CH^1_{AdS} which is periodic in t with period $\frac{1}{\omega}$ for some non-zero $\omega \in \mathbb{R}$, we have the expansion $\psi = \sum_{n=-\infty}^{\infty} e^{i\omega_n t}\hat{\psi}_n$, with $\omega_n = 2\pi \cdot n \cdot \omega$. By Theorem 1.3 all the $\hat{\psi}_n$ are zero.

1.3 Quasi-normal modes

The question of decay for solutions of wave equations in the presence of trapping has a long tradition in the context of the so-called *obstacle problem* in Minkowski space. Here, one studies the propagation of waves outside of a compact object. The analogue of trapped null-geodesics are piecewise straight lines reflected off the boundary of the obstacle but (due to the geometry of the obstacle) remaining in a compact region of space. For this problem, one can prove that, as in the setting studied here, a logarithmic local energy decay estimate always holds, even in the presence of stable trapping. In [9], this decay result was obtained via an analysis of the so-called *resonances*, also known as quasi-normal modes in the physics literature. Resonances are the poles of the meromorphic continuation of the (truncated) resolvent of the associated elliptic operator. In the black hole setting, such techniques have been very successfully applied on asymptoticall y de Sitter manifolds [32, 7, 35, 36, 22, 21].

Finally, the issue of locating the quasi-normal modes is also a popular theme in the physics literature [30, 23]. For the case of the wave equation on Schwarzschild-AdS, it was found numerically in [23] that the imaginary part of the quasi-normal modes approach the axis exponentially fast as the angular momentum mode ℓ goes to infinity. This is suggestive of general solutions decaying at best logarithmically.

1.4 The non-linear stability problem

Let us relate our results to the problem of non-linear stability for asymptotically AdS spacetimes. As mentioned above, there is no decay for the wave equation on pure AdS, in fact, one can construct an infinity of periodic solutions on this background. For the associated non-linear problem, that is the evolution under the Einstein equations of perturbations of Cauchy data for AdS (supplemented with appropriate boundary conditions), this is highly suggestive of instability. This fact was first pointed out in a talk of M. Dafermos [13], and suggested independently (using different arguments) by the results of [1]. Recently, there has been some interesting numerical work [6] in support of AdS being unstable.

Turning from pure AdS to black hole spacetimes, one has the result of [29] proving asymptotic stability of Schwarzschild-AdS with respect to *spherically* *symmetric* perturbations in the context of the Einstein-Klein-Gordon system. With this in mind, one may expect a similar result to hold for general perturbations of Schwarzschild-AdS, or more generally, Kerr-AdS spacetimes. On the contrary, the analysis of this paper suggests that this belief is likely to be misconceived: While the trapping mechanism is absent in the spherically symmetric case (allowing one to prove exponential decay in this symmetry class), it will, in general, prevent the linear solution from decaying faster than logarithmically. For the non-linear problem, the logarithmic decay rate of the linear solution will be too weak to close a standard bootstrap argument. Thus one may conjecture:

Conjecture 1.4. *Kerr-AdS spacetimes are dynamically unstable for generic perturbations.*

1.5 Outline

In Section 2, we present the family of Kerr-AdS Lorentzian manifolds. Sections 3 and 4 contain the functional framework for the Klein-Gordon field as well as a review of previous material (well-posedness and boundedness results) required later in the paper. In section 5, we exploit the separation of variables for the wave equation (1) to localize the solutions in various frequency ranges associated to both the time and the angular variables. This localization allows to replace (1) by an ordinary differential equation with a (frequency dependent) potential. The relevant properties of this potential are also proven in this section.

Our decay result is based on an integrated decay estimate for the low frequency part, which is derived via the construction of microlocalized energy currents which generate positive spacetime terms (pioneered in the black hole context in [15, 18, 19]). These currents are introduced in Section 6 and applied in Sections 7 and 8. A key ingredient is the use of *exponential* weights in these currents, with the constant in the exponential depending on the frequency cut-off. In Section 9, we collect our frequency space estimates to obtain an integrated decay estimate on the low frequency part of the solution. The high frequency part of the solution is estimated in Section 10 using simply the boundedness statement: This estimate gains a smallness factor (after commutation with angular derivatives) from the fact that the solution is supported on high frequencies. Finally, we conclud e the proof of the main theorem in Section 11 by interpolating between the high and low frequency estimates.

A special case of Kerr-AdS spacetimes are the Schwarzschild-AdS spacetimes. In this case, we can improve the estimates for the low frequency part of the solution to establish exponential decay of the energy (with the constant depending on the frequency cut-off). This is done in Section 12. Section 13 contains the proof of Theorem 1.3. This section can in fact be read independently of Sections 7–12. Finally, Appendix A contains some standard estimates associated to the frequency decomposition.

2 The Kerr-AdS family of spacetimes

The following presentation of Kerr-AdS spacetimes is strongly inspired by the presentation of the Kerr spacetimes found in [18].

2.1 The fixed manifold with boundary $\mathcal{R}$

Let $\mathcal{R}$ denote the manifold with boundary

$$\mathcal{R} = [0, \infty) \times \mathbb{R} \times \mathbb{S}^2.$$

We define standard coordinates y^* for $\mathbb{R}^+$, t^* for $\mathbb{R}$ and (θ, ϕ) for $\mathbb{S}^2$. This defines a coordinate system on $\mathcal{R}$, which is global modulo the well-known degeneration of the spherical coordinates.

We define the event horizon $\mathcal{H}^+$ to be the boundary of $\mathcal{R}$

$$\mathcal{H}^+ = \partial\mathcal{R} = \{y^* = 0\}.$$

We denote by T and Φ the vector fields ∂_{t^*} and ∂_ϕ, where the latter is to be understood as the extension of the coordinate vector field to all of $\mathbb{S}^2$ in the usual way. Let ϕ_τ denote the one-parameter family of diffeomorphisms generated by T. The manifold $\mathcal{R}$ will coincide with the domain of outer communication of the black hole spacetimes including the future event horizon $\mathcal{H}^+$.

2.2 The parameter space

The Kerr-AdS family of spacetimes depends on three parameters (a, M, l), which are respectively the angular momentum per unit mass, the mass and a quantity related to the cosmological constant Λ via $\Lambda = -\frac{3}{l^2}$. The parameters are assumed to satisfy $M > 0$ as well as $|a| < l$ (the latter is needed to exclude naked singularities). Specifically, we define the parameter space $\mathcal{P}$ to be

$$\mathcal{P} := \{(a, M, l) \in \mathbb{R} \times (0, \infty)^2 : |a| < l\}.$$

For any $(a, M, l) \in \mathcal{P}$, denote by $r_+(a, M, l)$ the largest real root of the polynomial

$$\Delta_-(r) = \left(r^2 + a^2\right)\left(1 + \frac{r^2}{l^2}\right) - 2Mr. \tag{4}$$

2.3 Kerr-AdS-star coordinates

Let $r(a, M, l, y^*)$ be a smooth real valued function defined on $\mathcal{P} \times [0, \infty)$ such that $r_{|(a,M,l)\times[0,\infty)}$ is a diffeomorphism onto $[r_+(a, M, l), \infty)$ which coincides with y^* for $y^* > 3M$. For each parameters $(a, M, l) \in \mathcal{P}$, the collection $(t^*, r(a, M, l, y^*), \phi, \theta)$ determines a coordinate system on $\mathcal{R}$, global up to the degeneration of the spherical coordinates, which we call *Kerr-AdS-star* coordinates.

2.4 The tortoise coordinate r^*

Given parameters $(a, M, l) \in \mathcal{P}$, we define the coordinate r^* on $\{r > r_+\}$ as

$$\frac{dr^*}{dr} = \frac{r^2 + a^2}{\Delta_-(r)} \qquad , \qquad r^*(r = +\infty) = \pi/2, \tag{5}$$

with $\Delta_-(r)$ as in (4). Note that $r^* = -\infty$ at the horizon. We will use the notation $R^\star_\infty = \frac{\pi}{2}$.

We also define $R^\star_{-\infty}$ to be a some value of $r^\star$ close to the horizon, i.e. $R^\star_{-\infty}$ is a large negative number, and we will eventually consider the limit $R^\star_{-\infty} \to -\infty$ in our estimates.

2.5 Boyer-Lindquist coordinates

Finally, given parameters $(a, M, l) \in \mathcal{P}$, we define Boyer-Lindquist coordinates $(t, r, \theta, \tilde{\phi})$ from the Kerr-AdS-star coordinates as follows:

$$t = t^* - A(r) \qquad \text{and} \qquad \tilde{\phi} = \phi - B(r) \tag{6}$$

where

$$\frac{dA}{dr} = \frac{2Mr}{\Delta_- \left(1 + \frac{r^2}{l^2}\right)} \qquad \text{and} \qquad \frac{dB}{dr} = \frac{a\left(1 - \frac{a^2}{l^2}\right)}{\Delta_-} \tag{7}$$

and A and B vanishes at infinity.

The Boyer-Lindquist coordinate system only covers $\mathrm{int}(\mathcal{R})$, i.e. the set $\{r > r_+\}$, which is the exterior of the black hole excluding the event horizon. Boyer-Lindquist coordinates are of particular relevance because it is with respect to these coordinates that the wave equation separates.

2.6 The Kerr-AdS metric for fixed (a, M, l)

We may now introduce the Kerr-AdS metric as the unique smooth extension to $\mathcal{R}$ of the tensor given on the Boyer-Lindquist chart by:

$$\begin{aligned} g_{KAdS} = {} & \frac{\Sigma}{\Delta_-} dr^2 + \frac{\Sigma}{\Delta_\theta} d\theta^2 + \frac{\Delta_\theta \left(r^2 + a^2\right)^2 - \Delta_- a^2 \sin^2\theta}{\Xi^2 \Sigma} \sin^2\theta d\tilde{\phi}^2 \\ & -2 \frac{\Delta_\theta \left(r^2 + a^2\right) - \Delta_-}{\Xi \Sigma} a \sin^2\theta \, d\tilde{\phi} dt - \frac{\Delta_- - \Delta_\theta a^2 \sin^2\theta}{\Sigma} dt^2 \end{aligned} \tag{8}$$

where Δ_- is defined by (4) and

$$\Sigma = r^2 + a^2 \cos^2\theta, \qquad \Delta_\pm = \left(r^2 + a^2\right)\left(1 + \frac{r^2}{l^2}\right) \pm 2Mr \tag{9}$$

$$\Delta_\theta = 1 - \frac{a^2}{l^2} \cos^2\theta, \qquad \Xi = 1 - \frac{a^2}{l^2}\,. \tag{10}$$

Note the components of the inverse in these coordinates:

$$g^{tt} = -\frac{\left(r^2+a^2\right)^2}{\Sigma\Delta_-} + \frac{a^2\sin^2\theta}{\Sigma\Delta_\theta}\,, \tag{11}$$

$$g^{t\tilde{\phi}} = \frac{\Xi\left(r^2+a^2\right)}{\Delta_-\Sigma}a - \frac{\Xi}{\Delta_\theta\Sigma}a\,, \qquad g^{\tilde{\phi}\tilde{\phi}} = \frac{\Xi^2}{\Sigma\Delta_\theta}\frac{1}{\sin^2\theta} - \frac{\Xi^2 a^2}{\Sigma\Delta_-}\,, \tag{12}$$

$$g^{rr} = \frac{\Delta_-}{\Sigma}, \qquad g^{\theta\theta} = \frac{\Delta_\theta}{\Sigma}, \qquad \sqrt{g} = \frac{\Sigma}{\Xi}\sin\theta\,. \tag{13}$$

That the tensor (8) indeed extends to a smooth metric on $\mathcal{R}$ is clear from examining its form in Kerr-AdS-star coordinates for which we have:

$$\begin{aligned}
g_{\theta\theta} &= \frac{\Sigma}{\Delta_\theta} \quad , \quad g_{t^\star t^\star} = g_{tt} \quad , && (14)\\
g_{\phi\phi} &= g_{\tilde{\phi}\tilde{\phi}} \quad , \quad g_{t^\star\phi} = g_{t\tilde{\phi}} \quad , && (15)\\
g_{rr} &= \frac{1}{\Sigma\left(1+\frac{r^2}{l^2}\right)^2}\left(\Delta_+ - a^2\sin^2\theta\left(k_0 + \frac{\Sigma}{l^2}\right)\right) \quad , && (16)\\
g_{\phi r} &= -\frac{a\sin^2\theta}{\Xi\Sigma k_0}\left[\Xi\Sigma + 2Mr\right] \quad , && (17)\\
g_{t^\star r} &= \frac{1}{\Sigma k_0}\left(2Mr - \frac{a^2}{l^2}\sin^2\theta\Sigma\right)\,. && (18)
\end{aligned}$$

with $k_0 = 1 + \frac{r^2}{l^2}$. The inverse components are

$$\begin{aligned}
g^{t^\star t^\star} &= -\frac{\Xi\Delta_+ + a^2\sin^2\theta\left(-k_0\Xi + \frac{2Mr}{l^2}\right)}{k_0^2\Delta_\theta\Sigma}\,, && (19)\\
g^{\theta\theta} &= \frac{\Delta_\theta}{\Sigma} \quad , \quad g^{\phi\phi} = \frac{\Xi^2}{\Delta_\theta\Sigma\sin^2\theta} \quad , \quad g^{t^\star\phi} = \frac{a\Xi}{k_0 l^2\Delta_\theta}\,, && (20)\\
g^{rr} &= \frac{\Delta_-}{\Sigma} \quad , \quad g^{\phi r} = \frac{a\Xi}{\Sigma} \quad , \quad g^{t^\star r} = \frac{2Mr}{k_0\Sigma}\,. && (21)
\end{aligned}$$

The unit normal of a constant $t^\star$ slice is

$$n_\Sigma = \sqrt{-g^{t^\star t^\star}}\partial_{t^\star} - \frac{g^{t^\star r}}{\sqrt{-g^{t^\star t^\star}}}\partial_r - \frac{g^{t^\star\phi}}{\sqrt{-g^{t^\star t^\star}}}\partial_\phi \quad , \; g(n_\Sigma, n_\Sigma) = -1\,, \tag{22}$$

with the determinant of the metric induced on constant $t^\star$ slices being

$$\sqrt{\det h} = \sqrt{\det g_{t^\star=const}} = \frac{\Sigma}{\Xi}\sin\theta\left(\sqrt{-g^{t^\star t^\star}}\right)\,. \tag{23}$$

2.7 Coordinate Vectorfields and Isometries

Note the following relation between the coordinate vectorfields in regular coordinates $(t^\star, r, \theta, \phi)$ and in Boyer-Lindquist tortoise coordinates $\left(t, r^\star, \theta, \tilde{\phi}\right)$:

$$\partial_{t^\star} = \partial_t \quad , \quad \partial_\phi = \partial_{\tilde{\phi}} \, ,$$

$$\partial_r = \frac{1}{1+\frac{r^2}{l^2}} \partial_t - \frac{r-r_+}{\Delta_-} \cdot \frac{a\Xi \left(r+r_+\right)}{r_+^2+a^2} \partial_{\tilde{\phi}} + \frac{r^2+a^2}{\Delta_-} \left(-\partial_t + \partial_{r^\star} - \frac{a\Xi}{r_+^2+a^2} \partial_{\tilde{\phi}} \right) .$$

This implies in particular that the vectorfield

$$Z = \frac{1}{\Delta_-} \left(-\partial_t + \partial_{r^\star} - \frac{a\Xi}{r_+^2+a^2} \partial_{\tilde{\phi}} \right) \tag{24}$$

extends continuously to the event horizon and that $\Delta_- Z\psi$ vanishes linearly in $r - r_+$ on the event horizon.

The vectorfields $T = \partial_t$ and $\Phi = \partial_{\tilde{\phi}}$ are both Killing fields. An important linear combination is the so-called Hawking-Reall vectorfield

$$K = T + \frac{a\Xi}{r_+^2+a^2} \Phi \, , \tag{25}$$

which in addition to being Killing is globally causal on the black hole exterior, provided that $|a|l < r_+^2$ holds. Note that K becomes null on the event horizon, while it is strictly timelike on the exterior.

2.8 Foliations

We denote by $\Sigma_{t^\star}$ (respectively Σ_t) the hypersurfaces of constant $t^\star$ (respectively constant t), cf. Figure 1.1 in the introduction. We write Σ_0 for the slice $\Sigma_{t^\star=0}$, which is where (for convenience) we will prescribe data for (1). We also make the convention that the Greek letter τ is always associated with the $t^\star$ foliation.

3 The norms

Let $\slashed{g}$ and $\slashed{\nabla}$ denote the induced metric and the covariant derivative on the spheres $\mathbb{S}^2_{t^\star,r}$ of constant $t^\star$ and r in $\mathcal{R}$.
We write $|\slashed{\nabla}...\slashed{\nabla}\psi|^2 = \slashed{g}^{AA'}...\slashed{g}^{BB'}\slashed{\nabla}_A...\slashed{\nabla}_B\psi\slashed{\nabla}_{A'}...\slashed{\nabla}_{B'}\psi$ to denote the induced norm on the spheres. We define the following non-degenerate energy norms for

the scalar field ψ, cf. [27]:

$$\begin{aligned}
\|\psi\|^2_{H^{0,s}_{AdS}(\Sigma_{t^\star})} &= \int_{\Sigma_{t^\star}} r^s \psi^2 r^2 dr \sin\theta d\theta d\phi \\
\|\psi\|^2_{H^{1,s}_{AdS}(\Sigma_{t^\star})} &= \int_{\Sigma_{t^\star}} r^s \left[r^2 \left(\partial_r \psi\right)^2 + |\slashed{\nabla}\psi|^2 + \psi^2 \right] r^2 dr \sin\theta d\theta d\phi \\
\|\psi\|^2_{H^{2,s}_{AdS}(\Sigma_{t^\star})} &= \|\psi\|^2_{H^{1,s}_{AdS}(\Sigma_{t^\star})} \\
&\quad + \int_{\Sigma_{t^\star}} r^s \left[r^4 \left(\partial_r \partial_r \psi\right)^2 + r^2 |\slashed{\nabla}\partial_r \psi|^2 + |\slashed{\nabla}\slashed{\nabla}\psi|^2 \right] r^2 dr \sin\theta d\theta d\phi
\end{aligned}$$

Higher order norms may be defined similarly. We denote by $H^{k,s}_{AdS}(\Sigma_{t^*})$, the space of functions ψ such that $\nabla^i \psi \in L^2_{loc}(\Sigma_{t^*})$ for $i = 0, ..., k$ and $\|\psi\|^2_{H^{k,s}_{AdS}(\Sigma_{t^\star})} < \infty$. We denote by $CH^{k,s}_{AdS}$ the set of functions ψ defined on $\mathcal{R}$, such that

$$\psi \in \bigcap_{q=0,..,k} C^q \left(\mathbb{R}_{t^\star} ; H^{k-q, s_q}_{AdS}(\Sigma_{t^\star}) \right)$$

where $s_k = -2$, $s_{k-1} = 0$ and $s_j = s$ for $j = 0, ..., k-2$.

When $s = 0$, we will feel free to drop the s in the notation, i.e. $H^{k,0}_{AdS} := H^k_{AdS}$ and $CH^k_{AdS} := CH^{k,0}_{AdS}$.

Remark 3.1. *One can actually prove boundedness and decay for norms with stronger r-weights in the angular derivatives than given above by commuting with angular momentum operators near infinity in the estimates.*

We also define the densities

$$\begin{aligned}
e_1[\psi] &= \frac{1}{r^2}\left(\partial_{t^\star}\psi\right)^2 + r^2\left(\partial_r \psi\right)^2 + |\slashed{\nabla}\psi|^2 + \psi^2 , \\
e_2[\psi] &= e_1[\partial_{t^\star}\psi] + \frac{1}{r^2}\left(\partial_{t^\star}\partial_{t^\star}\psi\right)^2 + e_1[\psi] \\
&\quad + r^4\left(\partial_r\partial_r\psi\right)^2 + r^2|\slashed{\nabla}\partial_r\psi|^2 + |\slashed{\nabla}\slashed{\nabla}\psi|^2 , \qquad (26)
\end{aligned}$$

which the reader can convert to Boyer-Lindquist coordinates using the formulae of Section 2.7. Finally, let us define the initial energies

$$\begin{aligned}
E_1[\psi] &= \|\psi\|^2_{H^{1,0}_{AdS}(\Sigma_0)} + \|\partial_{t^\star}\psi\|^2_{H^{0,-2}_{AdS}(\Sigma_0)} < \infty , \text{ for all } \psi \in CH^1_{AdS} \text{ and} \\
E_2[\psi] &= \|\psi\|^2_{H^{2,0}_{AdS}(\Sigma_0)} + \|\partial_{t^\star}\psi\|^2_{H^{1,0}_{AdS}(\Sigma_0)} + \|\partial_{t^\star}\partial_{t^\star}\psi\|^2_{H^{0,-2}_{AdS}(\Sigma_0)} < \infty ,
\end{aligned}$$

for all $\psi \in CH^2_{AdS}$.

4 Well-posedness and boundedness statement

We review in this section previous results concerning Klein-Gordon fields in Kerr-AdS spacetimes which will be used later in this paper.

From Definition 5.1 of [27] we recall that an H^2*-initial data set* (u, v, w) for (1) on a slice $\Sigma_{t^\star}$ of constant $t^\star$, say $t^\star = 0$ for convenience, is determined by specifying freely a pair $v \in H^{1,0}_{AdS}(\Sigma_0)$, $w \in H^{0,-2}_{AdS}(\Sigma_0)$, which then uniquely determines a function $u \in H^{2,s}_{AdS}(\Sigma_0)$ (for any $s < \sqrt{9-4\alpha}$) by solving an elliptic equation. The following statement is proven in [27]:

Theorem 4.1 (Well-posedness in the energy class, [27])**.** *Fix a Kerr-AdS spacetime with parameters* (M, l, a) *and let* $\alpha < 9/4$. *Let* (u, v, w) *be an* H^2_{AdS}*-initial data set on the spacelike slice* Σ_0*. Then, there exists a unique solution* ψ *of* (1) *such that* $\psi|_{t^\star=0} = u$, $\partial_t \psi|_{t^\star=0} = v$ *and such that* ψ *belongs to the class of* CH^2_{AdS} *functions (unique within* CH^1_{AdS}*).*

As usual, higher regularity is propagated and one can obtain the analogous statement for H^k_{AdS}-norms with arbitrary k.

We collect below also the main vector-field estimates leading to the uniform boundedness statement for solutions to (1). To state them in a form useful for the remainder of the paper, we recall the notion of the energy momentum tensor associated with ψ satisfying (1),

$$\mathbb{T}_{\mu\nu}\left[\psi\right] = \partial_\mu \psi \partial_\nu \psi - \frac{1}{2} g_{\mu\nu} \left(g^{\gamma\delta} \partial_\gamma \psi \partial_\delta \psi - \frac{\alpha}{l^2} \phi^2 \right) . \tag{27}$$

For a given spacetime vectorfield X, we define the energy current

$$J^X_\mu \left[\psi\right] = \mathbb{T}_{\mu\nu}\left[\psi\right] X^\nu , \tag{28}$$

which satisfies the identity

$$\nabla^\mu \left(J^X_\mu \left[\psi\right] \right) = \nabla^\mu \left(\mathbb{T}_{\mu\nu}\left[\psi\right] X^\nu \right) = \mathbb{K}_{\mu\nu}\left[\psi\right] {}^{(X)}\pi^{\mu\nu} , \tag{29}$$

with ${}^{(X)}\pi^{\mu\nu} = \frac{1}{2}\left(\nabla^\mu X^\nu + \nabla^\nu X^\mu\right)$ being the deformation tensor of X. The latter vanishes for X a Killing field.

Theorem 4.2. *Given parameters* $M > 0$ *and* $l > 0$ *there exists an* $a_{max} > 0$ *such that for all Kerr-AdS spacetimes with parameters* M, l, a *with* $|a| < a_{max}$*, the following is true: Any solution* ψ *of (1) with* $\alpha < \frac{9}{4}$ *arising via Theorem 4.1 from initial data defined on* Σ_0 *satisfies*

- *Conservation law:*

$$\int_{\Sigma_{t^\star}} J^K_\mu \left[\psi\right] n^\mu + \int_{\mathcal{H}(t^\star, 0)} \left[K\left(\psi\right)\right]^2 = \int_{\Sigma_0} J^K_\mu \left[\psi\right] n^\mu . \tag{30}$$

- *Control of non-degenerate norms: There exists a globally timelike vectorfield* N *(with* $-g\left(N, N\right) > c > 0$*) on the black hole exterior with*[3]

$$\|\psi\|^2_{H^{1,0}_{AdS}(\Sigma_{t^\star})} + \|\partial_{t^\star}\psi\|^2_{H^{0,-2}_{AdS}(\Sigma_{t^\star})} \lesssim \int_{\Sigma_{t^\star}} J^N_\mu \left[\psi\right] n^\mu . \tag{31}$$

---

[3] Recall that if $0 < \alpha < \frac{9}{4}$, then $J^N_\mu\left[\psi\right] n^\mu$ is actually not positive definite. However, the integral over $t^\star$ slices of that expression is, cf. [26].

The analogous statement holds with N replaced by K on the right- and norms degenerating near the horizon on the left-hand side implying that all of the terms appearing in (30) are non-negative.

- *Uniform boundedness statement:*

$$\int_{\Sigma_{t^\star}} J^N_\mu [\psi] n^\mu \lesssim \int_{\Sigma_0} J^N_\mu [\psi] n^\mu \tag{32}$$

  for all $t^\star$ slices to the future of Σ_0. Similar estimates hold for higher order energies.

- *Redshift estimate: There exists $3M > r_1 > r_0 > r_+$ such that the estimate*

$$\begin{aligned}\int_{\mathcal{R}(\tau_1,\tau_2)\cap\{r\leq r_0\}} J^N_\mu [\psi] n^\mu \lesssim \int_{\mathcal{R}(\tau_1,\tau_2)\cap\{r_0\leq r\leq r_1\}} J^N_\mu [\psi] n^\mu \\ +C_{M,l,a,\alpha}\int_{\mathcal{R}(\tau_1,\tau_2)\cap\{r\leq r_1\}} \phi^2 + \int_{\Sigma_{\tau_1}} J^N_\mu [\psi] n^\mu\end{aligned} \tag{33}$$

  holds for any $\tau_2 \geq \tau_1 \geq 0$.

In [26], the estimate (33) was crucial to obtain the boundedness statement, as the spacetime terms on the right hand side were shown to be controlled by the spacetime integral of $J^K_\mu [\psi] n^\mu$. Here, we will use (33) in conjunction with an integrated decay estimate in the interior for a certain range of frequencies. See Proposition 9.4 in Section 9.5.

5 Preliminaries

In the remainder of the paper, we study the solution ψ arising from initial data prescribed on Σ_0 as described in Theorem 4.1. We denote by $C = C_{M,l,a,\alpha}$ a constant depending only on the fixed parameters in the problem. The symbols $\lesssim$ and $\gtrsim$ are used to state inequalities which hold modulo such constants.

5.1 Time cut-off

Let $\chi(z)$ be a smooth real cut-off function which is equal to zero for $z \leq 0$ and equal to 1 for $z \geq 1$. For any $\tau \in \mathbb{R}$, we define $\chi_\tau(t^\star) = \chi(t^\star)\chi(\tau - t^\star)$ and finally

$$\psi^\tau(t^\star, \cdot) = \chi_\tau(t^\star)\ \psi(t^\star, \cdot) \tag{34}$$

which is supported in $0 \leq t^\star \leq \tau$ only. If ψ is a solution of the wave equation (1), then ψ^τ satisfies the inhomogeneous equation

$$\Box_{g_{KAdS}}\psi^\tau = F := 2\nabla^\alpha\chi^\star\nabla_\alpha\psi + (\Box_{g_{KAdS}}\chi^\tau)\psi\,, \tag{35}$$

with F supported in the two strips

$$\{0 \leq t^\star \leq 1\} \cup \{\tau - 1 \leq t^\star \leq \tau\} . \tag{36}$$

Note that

$$r^2 |F|^2 \leq \frac{1}{r^2} \left(\partial_{t^\star} \psi\right)^2 + r^2 \left(\partial_r \psi\right)^2 + |\nabla\!\!\!/ \psi|^2 + \psi^2 , \tag{37}$$

and hence in view of the support of F for any τ,

$$\int_{\mathcal{R}(0,\tau)} \left(r^2 |F|^2\right) r^2 dt^\star dr d\theta d\phi \lesssim \|\psi\|^2_{H^1_{AdS}(\Sigma_0)} . \tag{38}$$

Note also that

$$\int_{\Sigma_{t^\star}} J^N_\mu \left[\psi^\tau\right] n^\mu \lesssim \int_{\Sigma_{t^\star}} J^N_\mu \left[\psi\right] n^\mu \leq \|\psi\|^2_{H^1_{AdS}(\Sigma_0)} \tag{39}$$

and

$$\int_{\mathcal{H}(t^\star,0)} \left[K\left(\psi^\tau\right)\right]^2 \leq \|\psi\|^2_{H^1_{AdS}(\Sigma_0)} . \tag{40}$$

To see the latter, convert the zeroth order error-term in the strips arising from the cut-off into a spacetime term using the radial derivative of ψ. Finally, use the boundedness statement.

In view of the time cut-off, for any $r > r_+$, ψ^τ is now Schwartz in the t variable (with value in $H^{1,s}_{AdS}$) and we can take the Fourier transform,

$$\psi^\tau\left(t, r, \theta, \tilde{\phi}\right) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\omega t} \widehat{\psi^\tau}\left(\omega, r, \theta, \tilde{\phi}\right) d\omega . \tag{41}$$

We would like to decompose $\widehat{\psi^\tau}$ further using the separability properties of the wave equation. In this process, we will encounter certain self-adjoint operators on $\mathbb{S}^2$ (whose eigenfunctions are related to the oblate spheroidal harmonics). We will study the spectral properties of these operators first.

5.2 The (modified) oblate spheroidal harmonics

Define the $L^2\left(\sin\theta d\theta d\tilde{\phi}\right)$-self-adjoint operator P acting on $H^1\left(\mathbb{S}^2\right)$-complex valued functions (see section 7 of [18] for a more detailed discussion) as

$$\begin{aligned} -P\left(\xi\right) f \quad &= \quad \frac{1}{\sin\theta} \partial_\theta \left(\Delta_\theta \sin\theta \partial_\theta f\right) + \frac{\Xi^2}{\Delta_\theta} \frac{1}{\sin^2\theta} \partial^2_{\tilde{\phi}} f \\ &\quad + \Xi \frac{\xi^2}{\Delta_\theta} \cos^2\theta f - 2i\xi \frac{\Xi}{\Delta_\theta} \frac{a^2}{l^2} \cos^2\theta \; \partial_{\tilde{\phi}} f \ , \end{aligned} \tag{42}$$

with $\xi = a\omega$. We also define the operator P_α, which is equal to

$$P_\alpha\left(\xi\right) = \begin{cases} P\left(\xi\right) + \frac{\alpha}{l^2} a^2 \sin^2\theta & \text{if } \alpha > 0 , \\ P\left(\xi\right) + \frac{|\alpha|}{l^2} a^2 \cos^2\theta & \text{if } \alpha \leq 0 . \end{cases} \tag{43}$$

For $l \to \infty$ the operator reduces to the oblate spheroidal Laplacian on $\mathbb{S}^2$ considered in [18]. If also $a = 0$ we retrieve the Laplacian on the round sphere. Standard elliptic theory justifies the definitions

$$\begin{aligned} P\left(\xi\right) &\text{ has eigenvalues } \tilde{\lambda}_{m\ell}\left(\xi\right) \text{ with eigenfunctions } \tilde{S}_{m\ell}\left(a\xi, \cos\theta\right) e^{im\tilde{\phi}}, \\ P_\alpha\left(\xi\right) &\text{ has eigenvalues } \lambda_{m\ell}\left(\xi\right) \text{ with eigenfunctions } S_{m\ell}\left(a\xi, \cos\theta\right) e^{im\tilde{\phi}}. \end{aligned} \tag{44}$$

To analyse the spectral properties of these operators, we introduce[4]

$$\omega_+ = \frac{ma\Xi}{r_+^2 + a^2} \quad , \quad \omega_r\left(r\right) = \omega_r = \frac{ma\Xi}{r^2 + a^2}\,. \tag{45}$$

Lemma 5.1 (Spectral properties of $P_{(\alpha)} + \xi^2$)**.** *Both $\tilde{\lambda}_{m\ell}$ and $\lambda_{m\ell}$ satisfy the estimates*

$$\lambda_{m\ell} + \xi^2 \geq \Xi^2 \cdot |m|\left(|m| + 1\right)\,, \tag{46}$$

$$\lambda_{m\ell} + \xi^2 \geq \Xi^2 \cdot |m|\left(|m| + 1\right) + \frac{m^2 \Xi^2 a^4}{\left(r_+^2 + a^2\right)^2} - C_{a,l}|m||\omega - \omega_+|\,. \tag{47}$$

Proof. We will establish the estimate for $\tilde{\lambda}_{m\ell}$ (i.e. the operator $P\left(\xi\right)$). The same proof goes through simply neglecting the additional non-negative term present in $P_\alpha\left(\xi\right)$. Recalling (44) and writing shorthand $f = \tilde{S}_{m\ell}\left(\xi, \cos\theta\right) e^{im\tilde{\phi}}$, we find

$$\begin{aligned} &P\left(\xi\right) f + \xi^2 f = \\ -\frac{1}{\sin\theta}\partial_\theta\left(\Delta_\theta \sin\theta \partial_\theta f\right) - \frac{\Xi^2}{\Delta_\theta}\frac{1}{\sin^2\theta}\partial_{\tilde{\phi}}^2 f &+ \frac{\xi^2}{\Delta_\theta}\sin^2\theta f - 2m\xi\frac{\Xi}{\Delta_\theta}\frac{a^2}{l^2}\cos^2\theta f \\ = -\frac{1}{\sin\theta}\partial_\theta\left(\Delta_\theta \sin\theta \partial_\theta f\right) - \Xi^2 \frac{1}{\sin^2\theta}\partial_{\tilde{\phi}}^2 f &+ P_c\left(f\right) = \tilde{P}\left(\xi\right) f + P_c\left(f\right)\,, \end{aligned}$$

where

$$P_c = \frac{1}{\Delta_\theta}\left(\xi\sin\theta - m\Xi\frac{a^2}{l^2}\frac{\cos^2\theta}{\sin\theta}\right)^2 + \frac{\Xi^2 m^2 a^2}{l^2}\frac{\cos^2\theta}{\sin^2\theta} \geq 0\,. \tag{48}$$

For P_c one can also derive the estimate (use $\sin^{-2}\theta \geq 1 + \cos^2\theta$, $\Delta_\theta^{-1} \geq 1 + \frac{a^2}{l^2}\cos^2\theta$ and the $r_+^2 > |a|l$ condition)

$$P_c \geq \frac{m^2 \Xi^2 a^4}{\left(r_+^2 + a^2\right)^2} - C_{a,l}|m||\omega - \omega_+|\,. \tag{49}$$

---

[4] Recall that in the asymptotically flat case, the range $0 \leq m\omega \leq m\omega_+$ corresponds to the superradiant frequencies. In our case, there is no "superradiance" in view of the properties of the Hawking-Real vector field. Nevertheless it will be useful to rewrite the potential in terms of $\omega - \omega_+$, see below.

For $\tilde{P}$ we integrate by parts to see that

$$\int_{\mathbb{S}^2} \tilde{P}(\xi) f \cdot \bar{f} = \int_{\mathbb{S}^2} \left[\Delta_\theta |\partial_\theta f|^2 + \Xi^2 \frac{1}{\sin\theta} |\partial_{\tilde{\phi}} f|^2 \right] d\theta d\tilde{\phi} ,$$
$$\geq \Xi^2 \int_{\mathbb{S}^2} \left[|\partial_\theta f|^2 + \frac{1}{\sin^2\theta} |\partial_{\tilde{\phi}} f|^2 \right] \sin\theta d\theta d\tilde{\phi}$$

holds for any $f \in H^1\left(\mathbb{S}^2\right)$. □

5.3 The separation of variables

We are ready to obtain the fully separated wave equation. We decompose the $\widehat{\psi^\tau}$ of equation (41) as

$$\widehat{\psi^\tau}\left(\omega, r, \theta, \tilde{\phi}\right) = \sum_{m\ell} \left(\widehat{\psi^\tau}\right)^{(a\omega)}_{m\ell}(r)\, S_{m\ell}(a\omega, \cos\theta)\, e^{im\tilde{\phi}} \tag{50}$$

where ($d\tilde{\sigma} = \sin\theta d\theta d\tilde{\phi}$)

$$\left(\widehat{\psi^\tau}\right)^{(a\omega)}_{m\ell}(r) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2(t,r)} d\tilde{\sigma} e^{i\omega t}\ S_{m\ell}(a\omega, \cos\theta)\, e^{-im\tilde{\phi}}\ \psi^\tau\left(t, r, \theta, \tilde{\phi}\right).$$

The analogous Fourier-decomposition for the source term F in (35) yields

$$F^{(a\omega)}_{m\ell}(r) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2(t,r)} d\tilde{\sigma}\ e^{i\omega t}\ S_{m\ell}(a\omega, \cos\theta)\, e^{-im\tilde{\phi}}\ F\left(t, r, \theta, \tilde{\phi}\right).$$

After the renormalization

$$u^{(a\omega)}_{m\ell}(r) = \left(r^2 + a^2\right)^{\frac{1}{2}} \Psi^{(a\omega)}_{m\ell}(r) , \tag{51}$$

we finally obtain the equation

$$\left[u^{(a\omega)}_{m\ell}(r)\right]'' + \left(\omega^2 - V^{(a\omega)}_{m\ell}(r)\right) u = H^{(a\omega)}_{m\ell}(r) , \tag{52}$$

with the right hand side being equal to

$$H^{(a\omega)}_{m\ell}(r) = \frac{\Delta_-}{\sqrt{r^2 + a^2}} F^{(a\omega)}_{m\ell}(r) . \tag{53}$$

The potential $V^{(a\omega)}_{m\ell}(r)$ is defined as

$$V^{(a\omega)}_{m\ell}(r) = V^{(a\omega)}_{+,m\ell}(r) + V^{(a\omega)}_{0,m\ell}(r) + V_\alpha(r) , \tag{54}$$

where

$$\begin{aligned}
V^{(a\omega)}_{+,m\ell}(r) &= -\Delta_-^2 \frac{3r^2}{\left(r^2+a^2\right)^4} + \Delta_- \frac{5\frac{r^4}{l^2} + 3r^2\left(1+\frac{a^2}{l^2}\right) - 4Mr + a^2}{\left(r^2+a^2\right)^3} \\
&= \left(r^2+a^2\right)^{-\frac{1}{2}} \left(\sqrt{r^2+a^2}\right)'' , \qquad (55) \\
V^{(a\omega)}_{0,m\ell}(r) &= \frac{\Delta_-\left(\lambda_{m\ell}+\omega^2a^2\right) - \Xi^2a^2m^2 - 2m\omega a\Xi\left(\Delta_- - \left(r^2+a^2\right)\right)}{\left(r^2+a^2\right)^2} \\
V_\alpha(r) &= -\frac{\alpha}{l^2}\frac{\Delta_-}{\left(r^2+a^2\right)^2}\left(r^2+\Theta(\alpha)\,a^2\right). , \qquad (56)
\end{aligned}$$

with $\Theta(\alpha) = 1$ for $\alpha > 0$ and $\Theta(\alpha) = 0$ if $\alpha \leq 0$ (recall that the $\lambda_{m\ell}$ also depend on α from (43)). Note that V_+ grows like $\frac{2r^2}{l^4}$ near infinity while the V_0-part remains bounded.

5.4 Properties of the Potential

We will now study the potential in some detail. We will establish good positivity properties of the potential near the event horizon for all cases of Theorem 1.1. For $\alpha < 1$ we in fact obtain *global* (in r) positivity estimates. For the remaining α-range, we will have to allow for a negative contribution in the global estimate.

We will suppress the sub- and superscripts for the potential and its components in this subsection, i.e. we will write V for $V^{(a\omega)}_{m\ell}$, V_+ for $V^{(a\omega)}_{+,m\ell}$, etc.

Lemma 5.2 (Frequency independent properties)**.** *If any of the three conditions of Theorem 1.1 holds, there exists $\rho_+ > r_+$, depending only on a, l, M and α such that, for $r \in [r_+, \rho_+]$,*

$$V_+ + V_\alpha > C\frac{\Delta_-}{\left(r^2+a^2\right)^2}\, r^2, \qquad (57)$$

where C is a constant depending only on M, l, a, α. Moreover, again for any of the three conditions of Theorem 1.1, we have the global estimate

$$V_+ + V_\alpha + \frac{1}{4l^2}\frac{\Delta_-}{r^2+a^2} \geq C\frac{\Delta_-}{\left(r^2+a^2\right)^2}\, r^2 . \qquad (58)$$

For $\alpha < 1$ the last term on the left hand side of (58) can be dropped.

Proof. We have

$$V_+ = \frac{2\Delta_-}{\left(r^2+a^2\right)^2}\frac{r^2}{l^2} + \frac{\Delta_-}{\left(r^2+a^2\right)^4}\left(a^2\Delta_- + \left(r^2-a^2\right)2Mr\right) \geq 0 , \qquad (59)$$

from which one easily obtains (57) globally if $\alpha < 1$. For $\alpha < \frac{9}{4}$ it is straightforward to check that, adding the extra term on the left, (58) indeed holds under any of the conditions of Theorem 1.1. To obtain the estimate locally near the horizon without the extra term, we collect the leading order terms near the horizon from (55) to find that we need to establish positivity for

$$5\frac{r^4}{l^2}+3r^2\left(1+\frac{a^2}{l^2}\right)-4Mr+a^2-\frac{\alpha}{l^2}(r^2+a^2)^2 \tag{60}$$

at r_+. Using $\Delta_-(r_+)=0$ to evaluate the term $4Mr_+$, the sign of the above expression is the same as that of:

$$3\frac{r_+^4}{l^2}+r_+^2\left(1+\frac{a^2}{l^2}\right)-a^2-\alpha\frac{r_+^4}{l^2}-2\alpha\frac{r_+^2}{l^2}a^2-\alpha\frac{a^4}{l^2}, \tag{61}$$

which is easily seen to be strictly positive under any of the three conditions of Theorem 1.1. □

Note that one can obtain the estimate (57) near the horizon also for $(V_+ + V_\alpha)'$. For V_0, we simply remark the following property.

Lemma 5.3. *Setting $\mathcal{L}=\lambda_{m\ell}+\omega^2a^2$, we have*

$$V_0-\omega^2=-\left(\omega-\omega_r\right)^2+\frac{\Delta_-}{\left(r^2+a^2\right)^2}\left(\mathcal{L}-2ma\Xi\omega\right) \tag{62}$$

with the second term in (62) being non-negative.

Proof. The equality in (62) is an easy computation while the positivity claim is a consequence of the positivity of the operator $P+\xi^2-2m\xi\Xi$, which can be read off from

$$\left(P+\xi^2-2m\xi\Xi\right)f=-\frac{1}{\sin\theta}\partial_\theta\left(\Delta_\theta\sin\theta\partial_\theta f\right)+\frac{1}{\Delta_\theta}\left(\xi\sin\theta-\frac{m\Xi}{\sin\theta}\right)^2 f\,. \tag{63}$$

□

To perform further analysis near the horizon we introduce

$$V_{red}=V+|\omega-\omega_r(r)|^2-\omega^2\;,\qquad \tilde{V}=V+|\omega-\omega_+|^2-\omega^2\,. \tag{64}$$

From the previous Lemmata, we deduce

Lemma 5.4. *If any of the three conditions of Theorem 1.1 holds, there exists $\rho_+>r_+$, depending only on a, l, M and α, such that for all $r\in[r_+,\rho_+]$, we have:*

$$V_{red}\geq C\frac{\Delta_-}{\left(r^2+a^2\right)^2}\cdot r^2+\frac{\Delta_-}{(r^2+a^2)^2}\left(\mathcal{L}-2ma\Xi\omega\right) \tag{65}$$

with $C>0$ a constant depending only on a, l, M, α. Moreover, the estimate (65) is global for any of the conditions of Theorem 1.1, provided the term $\frac{1}{4l^2}\frac{\Delta_-}{r^2+a^2}$ is added to the left hand side when $\alpha\geq 1$.

In Section 7, we will analyze the behavior of solutions in the near stationary regime $\omega \sim \omega_+$. For this, we will need the following estimate on $\tilde{V}$,

Lemma 5.5. *Under any of the conditions of Theorem 1.1, we have the global estimate:*

$$\tilde{V} + \frac{1}{4l^2}\frac{\Delta_-}{r^2+a^2} \geq -C|m|\frac{\Delta_-}{\left(r^2+a^2\right)^2}|\omega-\omega_+| + C\frac{\Delta_-}{\left(r^2+a^2\right)^2}\left(m^2+r^2\right) \quad (66)$$

where the second term on the left hand side can be dropped if $\alpha < 1$. Moreover, if any of the three conditions of Theorem 1.1 holds, then there exists an $\Omega > 0$ and $\rho_+ > r_+$, depending only on a, l, M and α, such that for all frequencies $|\omega - \omega_+| \leq \frac{1}{\Omega}$ and all $r \in [r_+, \rho_+]$, we have

$$\tilde{V} \geq C\,\Delta_-\left[1+|m|\left(|m|+1\right)\right], \quad (67)$$

$$\tilde{V}' \geq C\,\Delta_-\left[1+|m|\left(|m|+1\right)\right], \quad (68)$$

$$\frac{\tilde{V}'}{\tilde{V}} \geq C\,. \quad (69)$$

with $C > 0$ a constant depending only on a, l, M and α.

Proof. We have

$$\tilde{V} = V_{red} + 2(\omega-\omega_+)(\omega_r-\omega_+) - (\omega_r-\omega_+)^2\,, \quad (70)$$

and using the properties of the $\lambda_{m\ell}$ collected in Lemma 5.1,

$$\begin{aligned}(\mathcal{L} - 2ma\Xi\omega) &= (\mathcal{L} - 2ma\Xi\omega_+) + 2ma\Xi(\omega-\omega_+) \\ &\geq \Xi^2|m|(|m|+1) + \frac{m^2a^4\Xi^2}{\left(r_+^2+a^2\right)^2} - 2ma\Xi\omega_+ - C_{a,l}|m||\omega-\omega_+|\,.\end{aligned}$$

We claim that

$$\left[\Xi^2m^2 - 2ma\Xi\omega_+ + \frac{m^2a^4\Xi^2}{\left(r_+^2+a^2\right)^2}\right]\frac{\Delta_-}{(r^2+a^2)^2} - (\omega_r-\omega_+)^2 \geq Cm^2\frac{\Delta_-}{\left(r^2+a^2\right)^2}$$

for some constant $C > 0$. To see this use the estimate

$$\frac{(r-r_+)(r+r_+)\left(r^2-r_+^2\right)}{\Delta_-} \leq l^2 \quad (71)$$

to find that the condition $r_+^2 > |a|l$ ensures the above estimate. This establishes (66). Assuming that $|\omega - \omega_+| \leq \frac{1}{\Omega}$ the estimates (67) and (68) immediately follow for Ω sufficiently large. For (69), recall the identity

$$\tilde{V} = V_+ + V_\alpha + \frac{\Delta_-}{\left(r^2+a^2\right)^2}\left(\mathcal{L} - 2ma\Xi\omega\right) + 2\left(\omega-\omega_+\right)\left(\omega_r-\omega_+\right) - \left(\omega_r-\omega_+\right)^2\,,$$

the fact that $\omega_r - \omega_+ \sim ma\Xi\Delta_-$ near the horizon and that $m^2 \lesssim \mathcal{L} - 2ma\Xi\omega_+$. From this we see that

$$\begin{aligned} \tilde{V} &\leq C_{M,l,a}\Delta_- \left(1 + (\mathcal{L} - 2ma\Xi\omega_+)\right) + C_{M,l,a}\left(\Omega^{-1}|m|\Delta_- + m^2\Delta_-^2\right) \\ &\leq C_{M,l,a}\Delta_- \left(1 + (\mathcal{L} - 2ma\Xi\omega_+)\right) \end{aligned} \tag{72}$$

holds for sufficiently large Ω and ρ_+ sufficiently close to the horizon. Similarly,

$$\begin{aligned} \tilde{V}' &\geq c_{M,l,a}\ \Delta_- \left(1 + (\mathcal{L} - 2ma\Xi\omega_+)\right) - C_{M,l,a}\Delta_- \left(\Omega^{-1}|m| + m^2\Delta_-\right) \\ &\geq c_{M,l,a}\ \Delta_- \left(1 + (\mathcal{L} - 2ma\Xi\omega_+)\right)\,, \end{aligned} \tag{73}$$

where we are exploiting both smallness of Ω^{-1} and of Δ_- to absorb the last term by the first. Combining (72) and (73) produces (69). $\square$

5.5 (ℓ, m, ω) low-frequency localisation

Let $\zeta(z)$ be a smooth cut-off function which is equal to 0 for $z \geq 2$ and equal to 1 for $z \leq 1$. Let moreover $\tilde{\zeta}(z)$ be a symmetric cut-off function being equal to 1 for $[-1/2, 1/2]$ and equal to zero for $|z| \geq 1$. We will cut off $\widehat{\psi^\tau}$ frequency both in the ω variable (at least in case that $a \neq 0$) and in the (ℓ, m) variables. Let us define, for fixed $L > 0$, $\Omega > 0$ the following kernels:

$$\begin{aligned} \mathcal{K}_{\leq L, \omega \not\approx \omega_+}\left(t - s, \theta, \underline{\theta}, \tilde{\phi}, \underline{\tilde{\phi}}\right) &= \int_{-\infty}^{\infty} d\omega \sum_{m,\ell} \zeta\left(\frac{a^2\omega^2 + \lambda_{m\ell}(\omega)}{L}\right) \cdot \\ \left[1 - \tilde{\zeta}\left((\omega - \omega_+)\Omega\right)\right] & e^{i\omega(t-s)} \cdot S_{m\ell}(a\omega, \cos\underline{\theta})\, S_{m\ell}(a\omega, \cos\theta)\, e^{im\left(\tilde{\phi} - \underline{\tilde{\phi}}\right)}\,, \end{aligned} \tag{74}$$

$$\begin{aligned} \mathcal{K}_{\leq L, \omega \approx \omega_+}\left(t - s, \theta, \underline{\theta}, \tilde{\phi}, \underline{\tilde{\phi}}\right) &= \int_{-\infty}^{\infty} d\omega \sum_{m,\ell} \zeta\left(\frac{a^2\omega^2 + \lambda_{m\ell}(\omega)}{L}\right) \cdot \\ \tilde{\zeta}\left((\omega - \omega_+)\Omega\right) e^{i\omega(t-s)} & \cdot S_{m\ell}(a\omega, \cos\underline{\theta})\, S_{m\ell}(a\omega, \cos\theta)\, e^{im\left(\tilde{\phi} - \underline{\tilde{\phi}}\right)}\,, \end{aligned} \tag{75}$$

and the associated projection operators

$$\begin{aligned} &\mathcal{P}_{\leq L, \omega \not\approx \omega_+}(\varphi)\left(t, r, \theta, \tilde{\phi}\right) = \\ &\int_{-\infty}^{\infty} ds \int_{\mathbb{S}^2} \sin\theta d\underline{\theta} d\underline{\tilde{\phi}}\ \varphi\left(s, r, \tilde{\theta}, \underline{\tilde{\phi}}\right) \mathcal{K}_{\leq L, \omega \not\approx \omega_+}\left(t - s, \theta, \underline{\theta}, \tilde{\phi}, \underline{\tilde{\phi}}\right)\,, \end{aligned}$$

for $\varphi\left(s, r, \underline{\theta}, \underline{\tilde{\phi}}\right)$ a smooth function. Similarly for $\mathcal{P}_{\leq L, \omega \approx \omega_+}(\varphi)\left(t, r, \theta, \tilde{\phi}\right)$. In applications, φ will be the cut-off scalar field ψ^τ or derivatives thereof. We finally define $\psi^\tau_\flat$, $\psi^\tau_\sharp$ as follows:

$$\begin{aligned} \psi^\tau_\flat &= \psi^\tau_{\flat, \omega \not\approx \omega_+} + \psi^\tau_{\flat, \omega \approx \omega_+} = \mathcal{P}_{\leq L, \omega \not\approx \omega_+}(\psi^\tau) + \mathcal{P}_{\leq L, \omega \approx \omega_+}(\psi^\tau)\,, \\ \psi^\tau_\sharp &= \psi^\tau - \psi^\tau_\flat\,. \end{aligned}$$

Naturally, one has $\psi^\tau_\flat$ and $\psi^\tau_\sharp$ in CH^k_{AdS} provided that ψ is in CH^k_{AdS}, for all $k \geq 1$.

6 Microlocal current templates

In this section, we shall define currents at the level of the separated Fourier components $u^{(a\omega)}_{m\ell}(r)$ introduced in section 5.3. These currents are clearly well-defined for $\psi^{\tau}_{\flat}$ since in this case the $u^{(a\omega)}_{m\ell}(r)$ are all smooth.

To avoid overloading the notation, we will employ in this section the short hand notation $u = u^{(a\omega)}_{m\ell}(r)$, $V = V^{(a\omega)}_{m\ell}(r)$ and $H = H^{(a\omega)}_{m\ell}(r)$.

6.1 Microlocal energy and red-shift currents

The microlocal current associated to the Killing fields $T = \partial_t$ and $K = \partial_t + \frac{a\Xi}{r_+^2+a^2}\partial_\phi$ is given by:

$$Q_T[u] = \omega Im\left(u'\bar{u}\right) \qquad \text{and} \qquad Q_K\left[u\right] = \left(\omega - \omega_+\right) Im\left(u'\bar{u}\right) \tag{76}$$

respectively. We have $Q'_T = \omega Im\left(H\overline{u}\right)$ and $Q'_K = \left(\omega - \omega_+\right) Im\left(H\bar{u}\right)$. Hence for $H = 0$, the quantities Q_T and Q_K are constant in $r^\star$.

Recall the definitions of (64). For any real function z, we also introduce the microlocal red-shift current:

$$Q^z_{red} = z\left[\left|u' + i\left(\omega - \omega_+\right)u\right|^2 - V_{red}\left|u\right|^2\right]. \tag{77}$$

The current Q^z_{red} satisfies

$$\begin{aligned} Q'_{red} =& z'\left|u' + i\left(\omega - \frac{a\Xi m}{r_+^2+a^2}\right)u\right|^2 - \left(zV_{red}\right)'\left|u\right|^2 \\ & - z(|u|^2)'\left(V_{red} - \tilde{V}\right) + 2zRe\left(\left(u' + i\left(\omega - \frac{a\Xi m}{r_+^2+a^2}\right)u\right)\overline{H}\right). \end{aligned} \tag{78}$$

6.2 Microlocal Morawetz currents

We finally turn to the Morawetz currents. Let f and g be bounded functions of $r^\star$ only. Using the notation of [18, 19], we define

$$Q^f_0 = f\left[|u'|^2 + \left(\omega^2 - V\right)|u|^2\right] + f'Re\left(u'\overline{u}\right) - \frac{1}{2}f''|u|^2, \tag{79}$$

$$Q^h_1 = hRe\left(u'\overline{u}\right) - \frac{1}{2}h'|u|^2, \tag{80}$$

$$Q^g_2 = g\left[|u'|^2 + \left(\omega^2 - V\right)|u|^2\right], \tag{81}$$

which satisfy

$$\left(Q^f_0\right)' = 2f'|u'|^2 - \left(fV' + \frac{1}{2}f'''\right)|u|^2 + 2fRe\left(u'\overline{H}\right) + f'Re\left(u\overline{H}\right), \tag{82}$$

$$\left(Q_1^h\right)' = h\left(|u'|^2 + \left(V-\omega^2\right)|u|^2\right) - \frac{1}{2}h''|u|^2 + hRe\left(u\overline{H}\right) \,, \tag{83}$$

$$\left(Q_2^g\right)' = g'\left[|u'|^2 + \left(\omega^2 - V\right)|u|^2\right] - gV'|u|^2 + 2gRe\left(u'\overline{H}\right) \,. \tag{84}$$

In the following, the combination

$$Q^{g,h} = Q_2^g + Q_1^h \tag{85}$$

will be particularly useful. For this choice

$$\begin{aligned}\left(Q^{g,h}\right)' &= |u'|^2\left(g'+h\right) + \omega^2\left(g'-h\right)|u|^2 \\ &+|u|^2\left(-g'V - gV' + Vh - \frac{1}{2}h''\right) + 2gRe\left(u'\overline{H}\right) + hRe\left(u\overline{H}\right) \,. \end{aligned} \tag{86}$$

7 Microlocal estimates for $\psi^\tau_{\flat,\omega\approx\omega_+}$

Recall the parameter Ω, which determines the scale where we cut-off near ω_+, cf. Section 5.5. In this section we prove an integrated decay estimate for frequencies very close to $\omega = \omega_+$. More precisely, we are going to determine an Ω (large) depending only on the parameters, such that for frequencies satisfying $|\omega - \omega_+| < \Omega^{-1}$, we have an integrated decay estimate. Once this constant Ω has been determined it will be fixed for the rest of the paper.

Remark 7.1. *These types of estimates for the "almost stationary regime" have appeared before in a different context in [18, 19] and recently in [3]. In fact, it is the latter version that we are adapting here because we want to avoid using the redshift at this point (cf. the remarks in Section 9.5).*

From Lemma 5.5, we first note that we can choose $\omega - \omega_+$ small enough such that we have the global estimate

$$\tilde{V} + \frac{1}{4l^2}\frac{\Delta_-}{r^2+a^2} \geq c_{M,l,a}\frac{\Delta_-}{\left(r^2+a^2\right)^2}\left(r^2+m^2\right) \,, \tag{87}$$

where the second term on the left hand side is not needed if $\alpha < 1$. For the mass range $\alpha \geq 1$ this additional term is gained from the derivative term via a Hardy inequality:

Lemma 7.2. *For any $r_{cut} \geq r_+$, we have for a smooth function u with $ur^{\frac{1}{2}} = o(1)$ at infinity*

$$\frac{1}{4l^2}\int_{r=r_{cut}}^\infty dr^\star \frac{\Delta_-}{\left(r^2+a^2\right)}|u|^2 \leq \int_{r=r_{cut}}^\infty dr^\star |u'|^2 \,. \tag{88}$$

Proof. Note that the weight on the left is the $r^\star$ derivative of r. Integrating by parts and Cauchy-Schwarz yields

$$\int_{r=r_{cut}}^\infty (r)'\,|u|^2\, dr^\star \leq 4\int_{r=r_{cut}}^\infty dr^\star \frac{r^2+a^2}{\Delta_-}\left(r-r_{cut}\right)^2 |u'|^2 \,. \tag{89}$$

The Lemma follows by estimating $r_{cut} > r_+$. $\square$

Next recall the multiplier (86) written in the form

$$\left(Q^{g,h}\right)' = |u'|^2 \left(g' + h\right) + g' \left(\omega - \omega_+\right)^2 |u|^2$$
$$+|u|^2 \left[\left(-g\tilde{V}\right)' - \frac{1}{2}h'' + h\left(V - \omega^2\right)\right] + 2gRe\left(u'\overline{H}\right) + hRe\left(u\overline{H}\right) . \qquad (90)$$

If $\omega = \omega_+$ we could simply choose $g = 0$ and $h = 1$ to obtain an integrated decay estimate (a variant of this argument will be used in Section 13 to exclude time-periodic solutions). However, as ω is only close to ω_+, we need to cut-off h and control the errors using the function g.

Let $r_+ < r_1 < r_2 < r_3$ (with corresponding $r^\star_{1,2,3}$) be r-values close to the horizon such that $r^\star_3 < 0$ and moreover (cf. Lemma 5.5)

$$\tilde{V} > 0\,, \quad \frac{\tilde{V}'}{\tilde{V}} \geq -\frac{2}{r^\star_2} \quad \text{and} \quad \frac{\tilde{V}'}{\tilde{V}} \geq \frac{1}{r}\frac{4\Delta_-}{r^2+a^2} \quad \text{hold in } r_+ < r \leq r_2\,.$$

From now on we regard r_2 and r_3 as being fixed, depending only on the above conditions and hence depending only on M, l and a and α in view of Lemma 5.5. The value of r_1 will be chosen below. Define $\underline{g}$ as follows:

$$\underline{g}\left(r^\star\right) = \begin{cases} \frac{1}{r^\star_1 \tilde{V}(r=r_1)} & \text{if } r < r_1, \\ \frac{1}{r^\star \cdot \tilde{V}(r)} & \text{if } r_1 \leq r < r_2, \\ \frac{r^\star - r^\star_3}{\left(r^\star_2 - r^\star_3\right) r^\star_2 \tilde{V}(r=r_2)} & \text{if } r_2 \leq r < r_3, \\ 0 & \text{if } r \geq r_3. \end{cases}$$

and hence

$$\underline{g}'\left(r^\star\right) = \begin{cases} 0 & \text{if } r < r_1, \\ -\frac{1}{r^\star \cdot \tilde{V}(r)}\left[\frac{\tilde{V}'}{\tilde{V}} + \frac{1}{r^\star}\right] & \text{if } r_1 \leq r < r_2, \\ \frac{1}{\left(r^\star_2 - r^\star_3\right) r^\star_2 \tilde{V}(r=r_2)} & \text{if } r_2 \leq r < r_3, \\ 0 & \text{if } r \geq r_3. \end{cases}$$

Note that $\underline{g}$ is C^0 and piecewise C^1 (hence the regularity is sufficient for the multiplier identity to hold), everywhere non-positive and monotonically increasing. Define $\underline{h}$ as

$$\underline{h}\left(r^\star\right) = \begin{cases} 0 & \text{if } r < r_1, \\ \tilde{h}\left(r^\star\right) & \text{if } r_1 \leq r < r_2, \\ \frac{4}{\left(r^\star_2 - r^\star_3\right) r^\star_2 \tilde{V}(r=r_2)} & \text{if } r \geq r_2. \end{cases}$$

where $\tilde{h}\left(r^\star\right)$ is a smooth monotone non-negative function interpolating smoothly between the two constant values at $r = r_1$ and $r = r_2$.

Using in particular that $\tilde{V}(r_2)$ can be bounded from below independently of m, we can choose r_1 sufficiently close to the horizon (depending only on M, l and a, in particular not on m) so that there exists a $\tilde{h}\left(r^\star\right)$ satisfying $|\tilde{h}''\left(r^\star\right)| < \frac{1}{(r^\star)^2}$ in $r_1 \leq r < r_2$. This fixes our $\tilde{h}\left(r^\star\right)$.

Finally, with Lemma 5.5 in mind, we choose Ω sufficiently large (again depending only on M, l and a) such that the following holds near the horizon for frequencies $|\omega - \omega_+| \leq \Omega^{-1}$:

$$
\begin{aligned}
V - \omega^2 &= \tilde{V} - \left(\omega - \omega_+\right)^2 \geq \frac{\tilde{V}}{2} \qquad \text{in } r_2 \leq r\,, \\
V - \omega^2 &= \tilde{V} - \left(\omega - \omega_+\right)^2 \geq 0 \qquad \text{in } r_1 \leq r \leq r_2\,.
\end{aligned}
$$

Note that with this choice we have in $r_1 \leq r \leq r_2$

$$
\underline{h}\left(V - \omega^2\right) + \left(-\underline{g}\tilde{V}\right)' - \frac{1}{2}\underline{h}'' \geq \frac{1}{\left(r^\star\right)^2} - \frac{1}{2\left(r^\star\right)^2} = \frac{1}{2\left(r^\star\right)^2}\,, \tag{91}
$$

while in $r_2 \leq r \leq r_3$ we have

$$
\begin{aligned}
\underline{h}\left(V - \omega^2\right) + \left(-\underline{g}\tilde{V}\right)' - \frac{1}{2}\underline{h}'' &= \underline{h}\left(\tilde{V} - \left(\omega - \omega_+\right)^2\right) - \underline{g}\tilde{V}' - \underline{g}'\tilde{V} + 0 \\
&\geq \left(\frac{\underline{h}}{2} - \underline{g}'\right)\tilde{V} \geq \frac{1}{L} c_{M,l,a} \frac{\Delta_-}{\left(r^2 + a^2\right)^2}\left(m^2 + r^2\right)\,,
\end{aligned}
\tag{92}
$$

with the L^{-1} appearing from the size of $\underline{h}$. Finally, for $r_3 \leq r < \infty$, $\underline{h}$ is constant. We can apply Lemma 7.2 with $r_{cut} = r_3$ to the derivative term $\underline{h}|u'|^2$ to gain the zeroth order term which makes (66) available. With this, we have, for sufficiently large Ω,

$$
\int_{r=r_3}^{r=\infty} dr^\star \ \underline{h}\left[\ |u'|^2 + \left(V - \omega^2\right)|u|^2\right] \geq \frac{c_{M,l,a}}{L} \int_{r_3}^{\infty} dr^\star \ |u|^2 \frac{\Delta_-}{\left(r^2 + a^2\right)^2}\left(m^2 + r^2\right).
$$

Note that this borrowing is not necessary if $\alpha < 1$. We summarize the construction in the following

Lemma 7.3. *Under any of the conditions in Theorem 1.1, there exists an Ω depending only on M, l and a such that the following holds for frequencies $|\omega - \omega_+| \leq \frac{1}{\Omega}$:*

$$
\begin{aligned}
&\frac{1}{L}\cdot c_{M,l,a}\left[\int_{r_2^\star}^{\infty} |u|^2 \frac{\Delta_-}{\left(r^2 + a^2\right)^2}\left(m^2 + r^2\right) dr^\star + \int_{r_1^\star}^{r_2^\star} \frac{1}{\left(r^\star\right)^2}|u|^2 dr^\star\right. \\
&\left. + \int_{R^\star_{-\infty}}^{r_1^\star} |u|^2 \frac{\Delta_-}{\left(r^2 + a^2\right)^2}\left(m^2 + r^2\right) dr^\star + \boxed{\int_{r_1^\star}^{\infty} |u'|^2 dr^\star}\ \right] \leq -\mathcal{Q}^{\underline{g},\underline{h}}\left(R^\star_{-\infty}\right) \\
&\qquad - \int_{R^\star_{-\infty}}^{\infty}\left[2\underline{g}Re\left(u'\overline{H}\right) + \underline{h}Re\left(u\overline{H}\right)\right] dr^\star
\end{aligned}
\tag{93}
$$

Here we have used that $\mathcal{Q}^{\underline{g},\underline{h}}(r^\star_\infty) = 0$ in view of the decay of u and u', cf. (151). The factor of L^{-1} appears since the functions $\underline{g}$ and $\underline{h}$ scale with L^{-1}. Note also that one first obtains this estimate without the boxed term, since it was used to borrow from in the Hardy inequality. However, one can simply repeat the estimate once more to regain control of the derivative term.

The $\underline{h}$-error-term in the last line of (93) is supported only for $r \geq r_1$ and easily absorbed on the left using Cauchy's inequality

$$|\underline{h} Re\left(u\overline{H}\right)| \leq |\underline{h}_{max}| \epsilon \frac{\Delta_-}{\left(r^2+a^2\right)^2}|u|^2 + \frac{|\underline{h}_{max}|}{\epsilon}\frac{\left(r^2+a^2\right)^2}{\Delta_-}|H|^2 \,. \tag{94}$$

As we will see later, the last term in (94) can be controlled from the boundedness statement. The $\underline{g}$-error in (93) is more problematic, since we do not have control over the derivative u' in the region $r \leq r_1$ on the left hand side. We need an auxiliary estimate. Choose $g = -r^{-2}$ and $h = 0$ in (90). Using again the properties of $\tilde{V}$ we obtain an estimate of the form

$$c_{M,l,a}\int_{R^\star_{-\infty}}^{r_3^\star}\left(|u'|^2 + \left(1+m^2\right)|u|^2\right)\frac{\Delta_-}{\left(r^2+a^2\right)^2}dr^\star \leq -\mathcal{Q}^{-r^{-1},0}\left(R^\star_{-\infty}\right)$$

$$\leq C_{M,l,a}\int_{r_3^\star}^{\infty}|u|^2\frac{\Delta_-^2}{\left(r^2+a^2\right)^4}\frac{1}{r}\left(L+r^2\right)dr^\star + C_{M,l,a}\int_{R^\star_{-\infty}}^{\infty}\frac{1}{r}|Re\left(u'\overline{H}\right)|dr^\star$$

Combining this estimate with the estimate of Proposition 7.3 and adding an angular term to the left hand side taking into account that $\lambda_{m\ell} + a^2\omega^2 \leq L$, we finally find

Proposition 7.4. *Under the condition of Lemma 7.3, we have*

$$\Bigg[\int_{R^\star_{-\infty}}^{\infty}\frac{\Delta_-}{\left(r^2+a^2\right)^2}\left[|u'|^2 + |u|^2\left(m^2+r^2+\lambda_{m\ell}+a^2\omega^2\right)\right]dr^\star \lesssim$$

$$-L^2\cdot\mathcal{Q}^{\underline{g},\underline{h}}\left(R^\star_{-\infty}\right) - L^2\cdot\mathcal{Q}^{-r^{-1},0}\left(R^\star_{-\infty}\right) + L^2\int_{R^\star_{-\infty}}^{\infty}dr^\star\frac{\left(r^2+a^2\right)^2}{\Delta_-}|H|^2\,, \tag{95}$$

where $u = \sqrt{r^2+a^2}\,\psi^{(a\omega)}_{m\ell}$ *denotes the Fourier-separated components of* $\psi^\tau_{\flat,\omega\approx\omega_+}$.

Remark 7.5. *The dependence on L could possibly be removed using a more refined analysis. However, since we are going to lose exponentially in L in the estimate for $\psi^\tau_{\flat,\omega\not\approx\omega_+}$, we content ourselves with this rather crude estimate.*

8 Microlocal estimates for $\psi^\tau_{\flat,\omega\not\approx\omega_+}$

In this section, we shall prove an integrated decay estimate for $\psi^\tau_{\flat,\omega\not\approx\omega_+}$. We will write Ψ for the separated Fourier-components of $\psi^\tau_{\flat,\omega\not\approx\omega_+}$ and as usual, $u = \sqrt{r^2+a^2}\,\Psi$.

To keep track of the various parameters involved in this section, we note already at this stage that we will, depending on the size of the mass α, first choose numbers n_2 and n_1. Finally, depending on Ω (fixed in the previous section), the constants n_1, n_2, the black hole parameters M, l, a and the cut-off constant L, we will choose a large constant $k = C_{M,l,a,\alpha,n_1,n_2} \cdot L \cdot \Omega^2$.

8.1 Preliminaries

Key to our argument in this frequency range is the use of exponentially weighted multipliers, which we introduce now.

8.1.1 Exponentially weighted multipliers

Let $x = \arctan\left(\frac{r}{l}\right)$. Note $c_{M,l,a} l \leq r^2 \cdot x_r = l\left[1 - \frac{1}{1+\frac{r^2}{l^2}}\right] \leq l$ and hence that the expression $r^2 x_r$ is essentially like a constant bounded away from zero and infinity. We define

$$f_n = \frac{1}{r^n} \exp\left(-k \arctan\left(\frac{r}{l}\right)\right) = \frac{1}{r^n} e^{-kx}\,, \tag{96}$$

$$f_n' = \left(\frac{-n}{r^{n+1}} - \frac{k\ x_r}{r^n}\right) \frac{\Delta_-}{r^2+a^2} e^{-kx} = \left(\frac{-n}{r} - k\ x_r\right) \frac{\Delta_-}{r^2+a^2} f_n = -\sigma f_n\,,$$

where

$$\sigma = \left(\frac{n}{r} + k\ \frac{(x_r r^2)}{r^2}\right) \frac{\Delta_-}{r^2+a^2}\,. \tag{97}$$

With this we compute

$$f_n'' = \left(\sigma^2 - \sigma'\right) f_n \qquad \text{and} \qquad f_n''' = -\left(\sigma^3 - 3\sigma\sigma' + \sigma''\right) f_n \tag{98}$$

Differentiating σ we find that away from the horizon, in $r \geq 3M$, we have for sufficiently large n and k (depending only on M, l, a) the bound

$$\sigma^3 - 3\sigma\sigma' + \sigma'' \geq c_{M,l,a}\ r^3 \left[n^3 + \frac{kn^2(x_r r^2)}{r} + \frac{nk^2(x_r r^2)^2}{r^2} + \frac{k^3(x_r r^2)^3}{r^3}\right] \tag{99}$$

To see this, note in particular that all terms on the left hand side grow like r^3 near infinity. For $r \leq 3M$ the expression on the left hand side of (99) is not positive everywhere. However, we can find a constant $C_{M,l,a}$ such that

$$\sigma^3 - 3\sigma\sigma' + \sigma'' + C_{M,l,a}\sigma \geq c_{M,l,a}\left(k^2\Delta_-^2 + k\Delta_-\right) \quad \text{holds in } r \leq 3M, \tag{100}$$

again provided k is chosen sufficiently large. We apply the same reasoning in the following Lemma, which has obviously been tailored to be applicable to the derivative of the multiplier (79) later.

Lemma 8.1. *If $f = f_n$ we have, for sufficiently large n and k (in particular $k \geq C_{M,l} L \geq C_{M,l}\left(\lambda_{m\ell} + a^2\omega^2\right)$),*

$$\begin{aligned} -\left(fV' + \frac{1}{2}f'''\right) \geq & -C_{M,l,a,\alpha} \cdot k \left(\frac{\Delta_-}{r^2+a^2}\right) \frac{e^{-kx}}{r^n}\chi_{\{r\leq 3M\}} \\ & + c_{M,l,a,\alpha}\left[k^2\left(\frac{\Delta_-}{r^2+a^2}\right)^2 \frac{1}{r^{n-1}}\right] e^{-kx} \end{aligned}$$

Remark 8.2. *Note that the first term on the right hand side is localized.*

Proof. Note the estimate

$$|fV'| \leq \frac{\Delta_-}{\left(r^2+a^2\right)^2} C_{M,l,a}\left(|\alpha| r^3 + Lr\right)\ \frac{e^{-kx}}{r^n} \qquad \text{for all } r \geq r_+ \tag{101}$$

Comparing with (99) and (100), it follows that this term can be absorbed by the f''' term for sufficiently large n and k. □

8.1.2 An exponentially weighted Hardy inequality

A second key tool is the use of Hardy inequalities with exponential weights.

Lemma 8.3. *Let $\beta \in [0,1]$ and $u(r^\star)$ be a smooth, bounded, complex valued function. Then, for positive numbers k and $n \geq 1$, we have the Hardy inequality*

$$\begin{aligned} & \int_{r=5M}^{\infty} \frac{e^{-kx}}{r^{n-1}}\left(n + \frac{k\left(x_r r^2\right)}{r}\right)|u|^2 dr^\star \\ \leq\ & 16 l^4 \int_{r=4M}^{\infty} \frac{e^{-kx}}{r^{n+1}}\left(\frac{1-\beta}{n} + \beta\frac{r}{k\left(x_r r^2\right)}\right)|u'|^2 dr^\star \\ & + C_{M,l,a}\int_{r=4M}^{r=5M} \frac{e^{-kx}}{r^{n-1}}\left(n + \frac{k\left(x_r r^2\right)}{r}\right)|u|^2 dr^\star . \end{aligned}$$

Proof. Let $\chi(r)$ be a cut-off function which is equal to 1 for $r \geq 5M$ and zero for $r \leq 4M$. Let $4M^\star$ denote the $r^\star$-value at $r = 4M$. We integrate by parts,

$$\begin{aligned} \int_{4M^\star}^{R^\star_\infty}\left(-\frac{e^{-kx}}{r^n}\right)'\chi|u|^2 dr^\star &= \int_{4M^\star}^{5M^\star}\frac{e^{-kx}}{r^n}\chi'|u|^2 dr^\star + 2\int_{4M^\star}^{R^\star_\infty}\frac{e^{-kx}}{r^n}\chi\ Re\left(\bar{u}u'\right) \\ &\leq \frac{1}{2}\int_{4M^\star}^{R^\star_\infty}\left(-\frac{e^{-kx}}{r^n}\right)'\chi\ |u|^2 dr^\star + 2\int_{4M^\star}^{R^\star_\infty}\frac{e^{-kx}e^{-kx}}{r^{2n}\left(-\frac{e^{-kx}}{r^n}\right)'}\ \chi\ |u'|^2 dr^\star \\ &\quad + \int_{4M^\star}^{5M^\star}\frac{e^{-kx}}{r^n}\chi'|u|^2 dr^\star \end{aligned}$$

observing that the boundary terms vanish in view of χ vanishing at $4M$ and u being bounded near infinity. This estimate leads to

$$\int_{4M^\star}^{R^\star_\infty} \left(-\frac{e^{-kx}}{r^n}\right)' \chi\, |u|^2 dr^\star$$

$$\leq 4\int_{4M^\star}^{R^\star_\infty} \frac{e^{-kx}e^{-kx}}{r^{2n}\left(-\frac{e^{-kx}}{r^n}\right)'}\,\chi\, |u'|^2 dr^\star + 2\int_{4M^\star}^{5M^\star} \frac{e^{-kx}}{r^n}\chi' |u|^2 dr^\star\,. \tag{102}$$

Observe now that

$$\left(-\frac{e^{-kx}}{r^n}\right)' = e^{-kx}\left(\frac{n}{r^{n+1}} + \frac{k\; r^2 \cdot x_r}{r^{n+2}}\right)\frac{\Delta_-}{r^2+a^2}\,. \tag{103}$$

Taking into account

$$\frac{1}{n+\frac{k(r^2 x_r)}{r}} \leq \frac{1-\beta}{n} + \beta\frac{r}{k\,(r^2 x_r)} \quad \text{and} \quad \frac{r^2+a^2}{\Delta} \leq \frac{2l^2}{r^2} \quad \text{for } r \geq 4M\,, \tag{104}$$

the inequality of the Lemma follows. $\square$

8.2 Estimating $|u'|^2$ from $|u|^2$

We turn to the estimates for the solution. We first apply the current Q_2^g with $g = -f_{n_2}$ (hence $g' = \sigma f_{n_2} > 0$). As in (90), we see that the spacetime term can be written

$$(Q_2^g)' = |u'|^2\; g' + |u|^2\left[g'\left(\omega-\omega_+\right)^2 + \left(-g\tilde{V}\right)'\right] + 2gRe\left(u'\overline{H}\right)\,. \tag{105}$$

We next study the behavior of the zeroth order term

$$\mathcal{V} = g'\left(\omega-\omega_+\right)^2 + \left(-g\tilde{V}\right)'\,. \tag{106}$$

Lemma 8.4. *In $r \leq 3M$, we have (with $g = -f_{n_2}$, $n_2 \geq 2$)*

$$-\frac{1}{2}g'\left(\omega-\omega_+\right)^2 + \left(g\tilde{V}\right)' \lesssim \Delta_-^2\; k\; \left(L+1\right)e^{-kx}\,. \tag{107}$$

Remark: The point here is that we have a *quadratic* degeneration near the horizon on the right hand side.

Proof. This estimate is easily seen for the term $g'\tilde{V}$ since it is already of order Δ_-^2. We will show that the Δ_--terms in the expression on the left hand side of (107) have a negative sign (all terms quadratic in Δ_- are again controlled by the right hand side in (107)). From Lemma 5.3, we recall

$$V - V_+ - V_\alpha - \omega^2 = -\left(\omega - \frac{ma\Xi}{r^2+a^2}\right)^2 + \frac{\Delta_-}{\left(r^2+a^2\right)^2}\left(\mathcal{L} - 2ma\Xi\omega\right) \tag{108}$$

and

$$\tilde{V}' = V' = (V_+ + V_\alpha)' + \frac{\Delta_-}{(r^2+a^2)^3}\Big[(\omega - \omega_r)\, ma\Xi \left(-2r - 2(\partial_r \Delta_-)\right)$$

$$+ (\partial_r \Delta_-)\left(\mathcal{L} - 2ma\Xi\omega_r\right)\Big] + \mathcal{O}\left(\Delta_-^2 L\right) .$$

The $(V_+ + V_\alpha)'$ term is already positive if terms of order Δ_-^2 are ignored. Note also that we can replace ω_r by ω_+ in the above since the difference $\omega_r - \omega_+ \sim m\Delta_-$ will have an additional degenerating factor near the horizon and produce terms of order $m^2\Delta_-^2 \leq L\Delta_-^2$. Ignoring the Δ_-^2 terms we need to show

$$-g\frac{\Delta_-}{(r^2+a^2)^3}\Big[\left(-2r - (2\partial_r \Delta_-)\right)(\omega - \omega_+)\, ma\Xi + (\partial_r \Delta_-)\left(\mathcal{L} - 2ma\Xi\omega_+\right)\Big]$$

$$+\frac{1}{2}g'\left(\omega - \omega_+\right)^2 \geq 0$$

holds in $r \leq 3M$. This is true in view of $\partial_r \Delta_-(r_+) > c_{M,l,a}$ and

$$\mathcal{L} - 2ma\Xi\omega_+ \geq \Xi^2 m^2 - 2m^2\frac{a^2}{r_+^2 + a^2}\Xi^2 \geq c_{M,l,a} \cdot m^2 \tag{109}$$

(ensuring a good m^2-term) and the fact that the $g'\left(\omega - \omega_+\right)^2$-term isf of size $k\left(\omega - \omega_+\right)^2$, which in combination with the good m^2-term will always allow us to absorb the cross-term for sufficiently large k. $\square$

Lemma 8.5. *In $r \geq 3M$, we have (with $g = -f_{n_2}$, $n_2 \geq 2$)*

$$-\frac{1}{2}g'\left(\omega - \omega_+\right)^2 + \left(g\tilde{V}\right)' \leq \frac{|2-\alpha|}{l^2}\left[\frac{n_2 - 2}{r^{n_2-3}}\frac{1}{l^2} + \frac{1}{l^2}\frac{k(x_r r^2)}{r^{n_2-2}}\right]e^{-kx}$$

$$+C_{M,l,a}\frac{k(x_r r^2)\left(L+1\right)}{r^{n_2-1}}e^{-kx} .$$

Proof. Near infinity/ in the interior we have (recall $\partial_{r^\star} \sim \frac{r^2}{l^2}\partial_r$ near infinity)

$$|\tilde{V} - \frac{2-\alpha}{l^2}\frac{r^2}{l^2}| \lesssim 1 + L + |m||\omega - \omega_+| \tag{110}$$

$$|\tilde{V}' - \frac{2\left(2-\alpha\right)}{l^2}\frac{r^3}{l^4}| \lesssim r + \frac{1}{r}L + \frac{1}{r}|m||\omega - \omega_+| \tag{111}$$

and hence

$$g\tilde{V}' + g'\tilde{V} \leq -f_{n_2}\left(\frac{2\left(2-\alpha\right)}{l^2}\frac{r^3}{l^4} - \frac{\Delta_-}{r^2+a^2}\left(\frac{n_2}{r} + \frac{k(x_r r^2)}{r^2}\right)\frac{2-\alpha}{l^2}\frac{r^2}{l^2}\right)$$

$$+C_{M,l,a,\alpha}\left[g'\left(1 + L + |m||\omega - \omega_+|\right) - g\left(r + \frac{1}{r}L + \frac{1}{r}|m||\omega - \omega_+|\right)\right]$$

$$\leq \frac{1}{2}g'\left(\omega - \omega_+\right)^2 + \frac{|2-\alpha|}{l^6}\left[\frac{n_2 - 2}{r^{n_2-3}} + \frac{k(x_r r^2)}{r^{n_2-2}}\right]e^{-kx}$$

$$+C_{M,l,a,\alpha}\left(n_2 + \frac{k(x_r r^2)}{r}\right)\frac{(L+1)}{r^{n_2-1}}e^{-kx}$$

since $m^2\Xi^2 < L$. $\square$

Combining the previous Lemmata, the estimate arising from the current Q_2^g with $g = -f_{n_2}$ finally reads

$$\begin{aligned}\int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \Bigg[\frac{\Delta_-}{\left(r^2+a^2\right)^2}\left[\left(\frac{n_2}{r^{n_2-1}} + \frac{k\ (r^2 x_r)}{r^{n_2}}\right)\right] e^{-kx}\left(|u'|^2\right) \\ + c_{M,l,a}\cdot k\left(\omega-\omega_+\right)^2 \frac{\Delta_-}{\left(r^2+a^2\right)^2} e^{-kx}\cdot\frac{1}{r^{n_2}}|u|^2\Bigg] \\ \leq \int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \Bigg(\frac{|2-\alpha|}{l^2}\left[\frac{n_2-2}{r^{n_2-3}} + \frac{k(x_r r^2)}{r^{n_2-2}}\right] e^{-kx} \\ + C_{M,l,a,\alpha}\left(n_2 + \frac{k(x_r r^2)}{r}\right)\frac{(L+1)}{r^{n_2-1}} e^{-kx}\Bigg)\frac{\Delta_-^2}{\left(r^2+a^2\right)^4}|u|^2 \\ + \left[Q_2^g\left(R^\star_\infty\right) - Q_2^g\left(R^\star_{-\infty}\right)\right] + \Big|\int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star\left[-2f_{n_2} Re\left(u'\overline{H}\right)\right]\Big| \end{aligned} \tag{112}$$

We claim that we can absorb (by the left hand side) the $|u|^2$-term on the right hand side in the region $r \geq 5M$ using Lemma 8.3. Indeed, for the first term on the right hand side of (112) we apply Lemma 8.3 with $n = n_2 - 2$ and $\beta = 0$: Choosing n_2 sufficiently large (depending only on the size of $|2-\alpha|$) then shows that this term is absorbed using a small part of the good $|u'|^2 \frac{n_2}{r^{n_2-1}}$ term on the left hand side. This also fixes our n_2. For the term in the second line (the one which has the constant $C_{M,l,a,\alpha}$), we apply Lemma 8.3 again, this time with the choice $n = n_2$ and $\beta = 1$. The Lemma reveals that for k sufficiently large, the term under consideration can be absorbed by the $|u'|^2 \frac{k(r^2 x_r)}{r^{n_2}}$ term on the left hand side. We summarize the new estimate as

$$\begin{aligned}\int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \Bigg[\frac{\Delta_-}{\left(r^2+a^2\right)^2}\left[\left(\frac{n_2}{r^{n_2-1}} + \frac{k\ (r^2 x_r)}{r^{n_2}}\right)\right] e^{-kx}\left(|u'|^2\right) \\ + c_{M,l,a}\cdot k\left(\omega-\omega_+\right)^2 \frac{\Delta_-}{\left(r^2+a^2\right)^2} e^{-kx}\cdot\frac{1}{r^{n_2}}|u|^2\Bigg] \\ \leq C_{M,l,a,\alpha}\int_{r\leq 5M} dr^\star\ \frac{k(x_r r^2)\,(L+1)}{r^{n_2-1}} e^{-kx}\frac{\Delta_-^2}{\left(r^2+a^2\right)^4}|u|^2 \\ + \left[Q_2^g\left(R^\star_\infty\right) - Q_2^g\left(R^\star_{-\infty}\right)\right] + \Big|\int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star\left[-2f_{n_2} Re\left(u'\overline{H}\right)\right]\Big|. \end{aligned} \tag{113}$$

In the following section, we prove a second multiplier estimate which essentially allows one to absorb also the part which is in $r \leq 5M$ by the left hand side.

8.3 Estimating $|u|^2$ from $|u'|^2$

We apply the current Q_0^f with $f = f_{n_1}$ and $n_1 \geq n_2 + 1$ sufficiently large so that Lemma 8.1 leads to the estimate

$$\begin{aligned}
\int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \left[\left(\frac{\Delta_-}{r^2+a^2} \right)^2 \frac{1}{r^{n_1-1}} \right] e^{-kx} |u|^2 \lesssim \frac{1}{k} \int_{r \leq 3M} dr^\star \left(\frac{\Delta_-}{r^2+a^2} \right) \frac{e^{-kx}}{r^{n_1-2}} |u|^2 \\
+ \frac{1}{k} \int_{-R^\star_{-\infty}}^{R^\star_\infty} dr^\star \left[\frac{\Delta_-}{\left(r^2+a^2\right)^2} \frac{1}{r^{n_1-1}} \right] e^{-kx} |u'|^2 + \frac{1}{k^2} \left[Q_0^{f_{n_1}} \left(R^\star_\infty\right) - Q_0^{f_{n_1}} \left(R^\star_{-\infty}\right) \right] \\
+ \frac{1}{k^2} \Big| \int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \left[2 f_{n_1} Re \left(u' \overline{H}\right) + f'_{n_1} Re \left(u \overline{H}\right) \right] \Big| .
\end{aligned} \tag{114}$$

8.4 Putting it together: The basic estimate

Estimating the term in $r \leq 5M$ on the right hand side of (113) by the estimate (114) and using that also $L \cdot \Omega^2 \ll k$, we arrive at

$$\begin{aligned}
\int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \frac{\Delta_-}{\left(r^2+a^2\right)^2} \frac{e^{-kx}}{r^{n_2}} \left(|u'|^2 + \left(\omega - \omega_+\right)^2 |u|^2 + \frac{1}{r} \frac{\Delta_-}{\left(r^2+a^2\right)^2} |u|^2 \right) \lesssim \\
\frac{1}{k^2} \left[Q_0^{f_{n_1}} \left(R^\star_\infty\right) - Q_0^{f_{n_1}} \left(R^\star_{-\infty}\right) \right] + \frac{1}{k^2} \Big| \int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \left[2 f_{n_1} Re \left(u' \overline{H}\right) + f'_{n_1} Re \left(u \overline{H}\right) \right] \Big| \\
+ \frac{1}{k} \left[Q_2^{-f_{n_2}} \left(R^\star_\infty\right) - Q_2^{-f_{n_2}} \left(R^\star_{-\infty}\right) \right] + \frac{1}{k} \Big| \int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \left[-2 f_{n_2} Re \left(u' \overline{H}\right) \right] \Big|
\end{aligned}$$

Multiplying by $e^{k\frac{\pi}{2}}$ and using elementary Hardy inequalities near the horizon and near infinity one can immediately improve the weights of the zeroth order term to

$$\begin{aligned}
\int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \frac{\Delta_-}{\left(r^2+a^2\right)^2} \frac{1}{r^{n_2}} \left(|u'|^2 + \left(\omega - \omega_+\right)^2 |u|^2 + r^2 |u|^2 \right) \lesssim \\
\frac{e^{k\frac{\pi}{2}}}{k^2} \left[Q_0^{f_{n_1}} \left(R^\star_\infty\right) - Q_0^{f_{n_1}} \left(R^\star_{-\infty}\right) \right] + \frac{e^{k\frac{\pi}{2}}}{k} \left[Q_2^{g} \left(R^\star_\infty\right) - Q_2^{g} \left(R^\star_{-\infty}\right) \right] \\
+ \frac{e^{k\frac{\pi}{2}}}{k^2} \Big| \int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \left[2 f_{n_1} Re \left(u' \overline{H}\right) + f'_{n_1} Re \left(u \overline{H}\right) \right] \Big| \\
+ \frac{e^{k\frac{\pi}{2}}}{k} \Big| \int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star \left[-2 f_{n_2} Re \left(u' \overline{H}\right) \right] \Big|
\end{aligned}$$

Finally, we can add an "angular" term, $k^{-1} \left(\lambda_{m\ell} + a^2 \omega^2\right) |\Psi|^2$ to the left hand side to obtain our basic integrated decay estimate.

Proposition 8.6. *For* $u=\sqrt{r^2+a^2}\psi_{m\ell}^{(a\omega)}$ *the Fourier-separated components of* $\psi^{\tau}_{\flat,\omega\not\approx\omega_+}$*, we have*

$$\begin{aligned}\int_{R^\star_{-\infty}}^{R^\star_{\infty}} dr^\star \frac{\Delta_- r^{-n_2}}{\left(r^2+a^2\right)^2}\left[|u'|^2+\left(\left(\omega-\omega_+\right)^2+r^2+r^2\left[\lambda_{m\ell}+a^2\omega^2\right]\right)|u|^2\right]\\ \lesssim \frac{e^{k\frac{\pi}{2}}}{k}\left[Q_0^{f_{n_1}}\left(R^\star_\infty\right)-Q_0^{f_{n_1}}\left(R^\star_{-\infty}\right)\right]+e^{k\frac{\pi}{2}}\left[Q_2^g\left(R^\star_\infty\right)-Q_2^g\left(R^\star_{-\infty}\right)\right]\\ +\frac{e^{k\frac{\pi}{2}}}{k}\Big|\int_{R^\star_{-\infty}}^{R^\star_{\infty}} dr^\star\left[2f_{n_1}Re\left(u'\overline{H}\right)+f'_{n_1}Re\left(u\overline{H}\right)\right]\Big|\\ +e^{k\frac{\pi}{2}}\Big|\int_{R^\star_{-\infty}}^{R^\star_{\infty}} dr^\star\left[-2f_{n_2}Re\left(u'\overline{H}\right)\right]\Big|. \qquad (115)\end{aligned}$$

9 Spacetime estimate for $\psi^{\tau}_{\flat}=\psi^{\tau}_{\flat,\omega\approx\omega_+}+\psi^{\tau}_{\flat,\omega\not\approx\omega_+}$

We succeeded in proving the microlocal estimates (95) and (115). We now want to turn these estimates into *spacetime estimates* by integration. For convenience, we will state the estimates in terms of $\psi=u\left(r^2+a^2\right)^{-1/2}$.

Proposition 9.1. *The low frequency part of* ψ^{τ}*,* $\psi^{\tau}_{\flat}$*, satisfies the integrated decay estimate*

$$\begin{aligned}\int_{\mathcal{D}}\left[\left(\partial_t\psi^{\tau}_{\flat}\right)^2+\left(\partial_{r^\star}\psi^{\tau}_{\flat}\right)^2+r^2|^{\gamma}\nabla\psi^{\tau}_{\flat}|^2_{\gamma}+r^2\psi^{\tau}_{\flat}\right]\frac{1}{r^{n_2+2}}dt^\star dr d\sigma_{\mathbb{S}^2}\\ \leq C_{M,l,a}\cdot e^{C_{M,l,a,\alpha}L}\int_{t^\star=0}J^N_\mu\left[\psi\right]n^\mu \qquad (116)\end{aligned}$$

where the integration is over the entire domain of outer communications, $\mathcal{D}$*.*

We will prove this estimate independently for $\psi^{\tau}_{\flat,\omega\approx\omega_+}$ (from (95)) and $\psi^{\tau}_{\flat}$ (from (115)), hence for the sum. Later, we will improve further both the weights near infinity (using an r^p-weighted estimate) and near the horizon (using the redshift) below.

Proof. Integrate (95) and (115) in ω and sum over l and m to convert the two estimates into spacetime integrated decay estimates for $\psi^{\tau}_{\flat,\omega\approx\omega_+}$ and $\psi^{\tau}_{\flat,\omega\not\approx\omega_+}$ respectively. The precise estimation of the terms which arise is contained in sections 9.1-9.3. ☐

9.1 The main terms

Replacing u by $\sqrt{r^2+a^2}\psi$, the term on the left hand side of (115) can be estimated

$$\lim_{R^\star_{-\infty}\to-\infty}\int_{-\infty}^{\infty} d\omega \int_{R^\star_{-\infty}}^{R^\star_{\infty}} dr^\star \sum_{m\ell}\left\{\frac{\Delta_-}{\left(r^2+a^2\right)^2}\frac{1}{r^{n_2-2}}\right.$$

$$\left.\left(\left|\left(\Psi^{(a\omega)}_{m\ell}\right)'\right|^2+\omega^2|\Psi^{(a\omega)}_{m\ell}|^2+r^2\left(\Psi^{(a\omega)}_{m\ell}\right)^2+r^2\left[\lambda_{m\ell}+a^2\omega^2\right]\left(\Psi^{(a\omega)}_{m\ell}\right)^2\right)\right\}$$

$$\gtrsim \int_{\mathcal{D}}\left[\left(\partial_t\psi^\tau_{\flat,\omega\not\approx\omega_+}\right)^2+\left(\partial_{r^\star}\psi^\tau_{\flat,\omega\not\approx\omega_+}\right)^2+r^2|{}^\gamma\nabla\psi^\tau_{\flat,\omega\not\approx\omega_+}|^2_\gamma\right.$$

$$\left.+r^2\left(\psi^\tau_{\flat,\omega\not\approx\omega_+}\right)^2\right]\frac{\Delta_-}{\left(r^2+a^2\right)^2}\frac{1}{r^{n_2-2}}dtdr^\star d\sigma_{\mathbb{S}^2}\,.$$

Here we have used the estimate (a consequence of Lemma 5.1)

$$\int_{-\infty}^{\infty}\sum_{m\ell}\left(\lambda_{m\ell}\left(a\omega\right)+a^2\omega^2\right)|\Psi^{(a\omega)}_{m\ell}|^2d\omega$$

$$\geq c_{M,l,a}\int_{\infty}^{\infty}dt\int d\sigma_{\mathbb{S}^2}|{}^\gamma\nabla\psi^\tau_\flat|^2_\gamma\,, \tag{117}$$

where γ denotes the standard unit metric on the (t,r)-spheres. The same estimate holds for the left hand side of (95) with $\psi^\tau_{\flat,\omega\approx\omega_+}$ replacing $\psi^\tau_{\flat,\omega\not\approx\omega_+}$ everywhere.

9.2 The H-error-terms

For the H-error-terms in (115) we estimate for a weight function $w\left(r\right)$

$$\lim_{R^\star_{-\infty}\to-\infty}\sum_{m,\ell}\int_{-\infty}^{\infty} d\omega \int_{R^\star_{-\infty}}^{R^\star_{\infty}} dr^\star\ w\left(r\right) Re\left(u'\overline{H}\right)=$$

$$\lim_{R^\star_{-\infty}\to-\infty}\sum_{m,\ell}\int_{-\infty}^{\infty} d\omega \int_{R^\star_{-\infty}}^{R^\star_{\infty}} dr^\star\ w\left(r\right) Re\left(\Psi'\sqrt{r^2+a^2}\,\overline{H}+\Psi\frac{r\Delta_-}{\left(r^2+a^2\right)^{\frac{3}{2}}}\overline{H}\right)$$

$$\leq \epsilon\cdot\lim_{R^\star_{-\infty}\to-\infty}\sum_{m,\ell}\int_{-\infty}^{\infty} d\omega \int_{R^\star_{-\infty}}^{R^\star_{\infty}} dr^\star\ \left(r^2|\Psi|^2+|\Psi'|^2\right)\frac{\Delta_-}{r^4}r^{-2p}$$

$$+\frac{C_{M,l}}{\epsilon}\cdot\lim_{R^\star_{-\infty}\to-\infty}\sum_{m,\ell}\int_{-\infty}^{\infty} d\omega \int_{R^\star_{-\infty}}^{R^\star_{\infty}} dr^\star\ |H|^2\left[r^{1+p}\cdot w\left(r\right)\right]^2\frac{r^4}{\Delta_-}$$

for any $\epsilon > 0$ and $p \geq 0$. For the last term we further estimate

$$\begin{aligned}
&\lim_{R^\star_{-\infty}\to-\infty}\sum_{m,\ell}\int_{-\infty}^{\infty}d\omega\int_{R^\star_{-\infty}}^{R^\star_{\infty}}dr^\star\ |H|^2\left[r^{1+p}\cdot w\left(r\right)\right]^2\frac{r^4}{\Delta_-}\\
\leq\ &\lim_{R^\star_{-\infty}\to-\infty}\sum_{m,\ell}\int_{-\infty}^{\infty}d\omega\int_{R^\star_{-\infty}}^{R^\star_{\infty}}dr^\star\ |F^{(a\omega)}_{m\ell}|^2\left[r^{1+p}\cdot w\left(r\right)\right]^2\Delta_-\cdot r^2\\
\leq\ &\lim_{R^\star_{-\infty}\to-\infty}\int_{-\infty}^{\infty}dt\int_{R^\star_{-\infty}}^{R^\star_{\infty}}dr^\star\int_{\mathbb{S}^2}r^2\sqrt{\gamma}d\theta d\tilde{\phi}\ |F_\flat|^2\left[r^{1+p}\cdot w\left(r\right)\right]^2\Delta_-\\
\leq\ &\lim_{R^\star_{-\infty}\to-\infty}\int_{-\infty}^{\infty}dt\int_{R^\star_{-\infty}}^{R^\star_{\infty}}dr^\star\int_{\mathbb{S}^2}r^2\sqrt{\gamma}d\theta d\tilde{\phi}\ r^2|F|^2\left[r^{1+p}\cdot w\left(r\right)\right]^2\frac{\Delta_-}{r^2}\\
&\lesssim \|\psi\|^2_{H^1_{AdS}(\Sigma_0)} \qquad (118)
\end{aligned}$$

provided $|r^{1+p}\cdot w\left(r\right)| \leq C$ is uniformly bounded. Here the last step follows from (38). All error-terms appearing in (115), namely

$$Re\left(u\overline{H}\right)f^\prime_{n_1} \quad , \quad Re\left(u^\prime\overline{H}\right)f_{n_1} \quad , \quad Re\left(u^\prime\overline{H}\right) - 2f_{n_2} \qquad (119)$$

can be treated in this way as is readily checked by inspecting the decay of the weights. Clearly, the H-error in (95) is also controlled by $\|\psi\|^2_{H^1_{AdS}(\Sigma_0)}$ in view of the previous computation.

9.3 The boundary-terms

First a rough outline of the argument: The boundary terms at infinity will be easily seen to vanish. The boundary term on $R^\star_{-\infty}$ becomes – after integration over ω, m, ℓ – a boundary term on a timelike hypersurface of constant $R^\star_{-\infty}$ on the black hole exterior. In the limit $R^\star_{-\infty} \to -\infty$ this term becomes a boundary integral on the horizon, which is in turn controlled by the energy of the Hawking-Reall Killing vectorfield.

We now turn to the precise structure of the terms. Note that all boundary terms at $R^\star_{-\infty}$ in both (115) and (95) are of the form

$$Q^{\cdots}\left(R^\star_{-\infty}\right) = y\left[|u^\prime|^2 + \left(\omega - \frac{ma\Xi}{r^2+a^2}\right)^2|u|^2\right] + \text{terms with } \Delta_-\text{-factor} \qquad (120)$$

for a bounded function $y = y\left(r^\star\right)$. Recall that the vectorfield Z defined in (24) is regular on the future event horizon. In particular, the quantities $Z\left(\psi^\tau\right)$, $Z\left(\psi^\tau_\flat\right)$ are bounded pointwise. In Fourier-space, this condition translates to the condition that

$$\lim_{r^*\to-\infty}\left[u^\prime + i\left(\omega - \frac{ma\Xi}{r^2+a^2}\right)u\right] = 0 \qquad (121)$$

holds. Recalling the flux of the Hawking Reall vectorfield through the horizon

$$\int_{\mathcal{H}^+} \left(\partial_t \psi + \frac{ma\Xi}{r_+^2 + a^2} \partial_\phi \psi \right)^2 \leq \int_{\mathcal{H}^+} J^K_\mu \left[\psi \right] n^\mu_{\mathcal{H}^+} \, , \tag{122}$$

one obtain the estimate ("+0" indicating terms vanishing in the limit)

$$\begin{aligned} & \lim_{R^\star_{-\infty} \to -\infty} \int_{R^\star_{-\infty}}^\infty d\omega \sum_{m\ell} \left\{ - \tilde{Q}^{\cdots} \left(R^\star_{-\infty} \right) \right\} \\ \lesssim & \lim_{R^\star_{-\infty} \to -\infty} \int_{-\infty}^\infty dt \int_{\mathbb{S}^2} \sin\theta d\theta d\tilde{\phi} \, | K \psi^\tau_\flat \left(t, r\left(R^\star_{-\infty} \right), \theta, \tilde{\phi} \right) |^2 + 0 \\ \leq & \lim_{R^\star_{-\infty} \to -\infty} \int_{-\infty}^\infty dt \int_{\mathbb{S}^2} \sin\theta d\theta d\tilde{\phi} \, | K \psi^\tau \left(t, r\left(R^\star_{-\infty} \right), \theta, \tilde{\phi} \right) |^2 + 0 \\ & = \int_{\mathcal{H}^+(0,\tau)} | K \left(\psi^\tau \right) |^2 \leq \| \psi \|^2_{H^1_{AdS}(\Sigma_0)} \, , \end{aligned} \tag{123}$$

with the last step following from (40).

9.4 Improving the weights near null-infinity

We can improve the weights near infinity in Proposition 9.1, so that we actually control the full energy in time (of course, it is still a degenerate energy near the horizon). This feature is characteristic of the asymptotically AdS end and was first observed in [29].

The following Hardy inequality will be useful, which can be proven using a simple integration by parts and Cauchy-Schwarz (or deduced from Lemma 7.2).

Lemma 9.2. *Given $\epsilon > 0$, there exists an $\tilde{R}^\star$ depending only on the parameters a, M, l such that*

$$\int_{\tilde{R}^\star}^{R^\star_\infty} dr^\star r^2 |u|^2 \leq \left[4 - \epsilon \right] l^4 \int_{\tilde{R}^\star}^{R^\star_\infty} dr^\star |u'|^2 \, . \tag{124}$$

Proposition 9.3. *We have the estimate*

$$\begin{aligned} \int_{\mathcal{D}} \left[\left(\partial_t \psi^\tau_\flat \right)^2 + r^2 \left(\partial_{r^\star} \psi^\tau_\flat \right)^2 + r^2 |^\gamma \nabla \psi^\tau_\flat |^2_\gamma + r^4 \psi^\tau_\flat \right] \frac{1}{r^2} dt^\star dr d\sigma_{\mathbb{S}^2} \\ \leq C_{M,l,a} \cdot e^{C_{M,l,a} L \cdot \Omega^2} \int_{t^\star = 0} J^N_\mu \left[\psi \right] n^\mu \, . \end{aligned} \tag{125}$$

Remark: Inspecting the proof of Proposition 9.1, we see that we already proved this for $\psi^\tau_{\flat, \omega \approx \omega_+}$.

Proof. Choose $f = -\frac{1}{r}$ for the microlocal current Q^f_0 in (79). Repeating the conversion to physical space of the previous section, one sees that the boundary terms near infinity and the horizon can be dealt with as before using only the boundedness statement. What does the resulting spacetime term control? Inspecting (82), we see that

$$\left(Q_0^f\right)' \geq \frac{2\Delta_-}{r^2\left(r^2+a^2\right)}|u'|^2 - C_{M,l,a}\left(1+\frac{L}{r}\right)|u|^2 + 2\left(2-\alpha\right)\frac{r^2}{l^6}|u|^2 - \text{H-Err}\,.$$

Let us choose an $r_L > r_+$ so large that both

$$\frac{\Delta_-}{r^2\left(r^2+a^2\right)} \geq \frac{1}{l^2} + \frac{1}{2r^2} \tag{126}$$

and also

$$\frac{1}{2}\left[\frac{1}{2} + 2\left(2-\alpha\right)\right]\frac{r^2}{l^6} \geq C_{M,l,a}\left(1+\frac{L}{r}\right) \tag{127}$$

hold for $r \geq r_L$. Note that (besides the dependence of M, l and a) r_L will depend polynomially (in fact, linearly) on L, and like $(9/4-\alpha)^{-1}$ on the mass. We now apply Lemma 9.2 (possibly making r_L a bit larger), which guarantees

$$\int_{r_L^\star}^{R_\infty^\star}\left(Q_0^f\right)' dr^\star \geq c_{M,l,a,\alpha}\int_{r_L^\star}^{R_\infty^\star} 2\frac{\Delta_-}{r^2}\left(|\Psi'|^2 + r^2\Psi^2\right)dr^\star - \text{H-Err}\,.$$

In the region $r \leq r_L$ we can estimate the r-weights by L-weights to find

$$\Big|\int_{R_{-\infty}^\star}^{r_L^\star}\left(Q_0^f\right)' dr^\star\Big| \lesssim L^{n_2}\int_{R_{-\infty}^\star}^{r_L^\star}\frac{2\Delta_-}{\left(r^2+a^2\right)^2}\frac{1}{r^{n_2-2}}\left(|\Psi'|^2 + r^2\Psi^2\right)dr^\star - \text{H-Err},$$

with the first term being already controlled from our basic estimate. The fact that we lose polynomially in L here is irrelevant in view of the exponential loss which has already occurred.

Finally, the H-error terms can be treated as previously, namely applying the estimates of Section 9.2 with $w\left(r\right) = r^{-1}$ and $p = 0$). Hence the estimate of the Proposition follows. $\square$

9.5 Improving the weights near the horizon

We would like to improve the weights near the horizon in Proposition 9.3 so that the left hand side is actually the non-degenerate energy integrated in time. However, we cannot construct a microlocal redshift estimate applying the techniques by which we derived the integrated decay estimate of Proposition 9.3. The reason is that the resulting boundary term will not have a good sign near the horizon (since it will be supported on both the future and the past horizon). One could of course estimate this boundary term by the (non-degenerate) N-energy of ψ^τ on the future event horizon. However, the latter actually diverges in the limit as $\tau \to \infty$, since the full solution ψ only decays logarithmically. This is in sharp contrast with the situation, when we derived the integrated decay estimate of Proposition 9.3: Here the boundary term could be localized on the future event horizon and controlled by the (degenerate!) K-energy of ψ^τ, which *is* controlled from the boundedness statement.[5]

The resolution is to apply the N-estimate in physical space locally between two spacelike $\Sigma_{t^\star}$-slices. The only problem, then, is to get a good bound for the N-energy of $\psi^\tau_\flat$ (from the total energy) on some initial slice. This will be achieved later using the pigeonhole principle. We start with the local N-estimate for $\psi^\tau_\flat$:

Proposition 9.4.

$$\int_{\Sigma_{\tau_2}} J^N_\mu \left[\psi^\tau_\flat\right] n^\mu + \int_{\mathcal{R}(\tau_1,\tau_2)} J^N_\mu \left[\psi^\tau_\flat\right] n^\mu \lesssim \int_{\Sigma_{\tau_1}} J^N_\mu \left[\psi^\tau_\flat\right] n^\mu + C_{M,l,a} \cdot e^{C_{M,l,a} L \frac{\pi}{2}} \int_{t^\star=0} J^N_\mu \left[\psi\right] n^\mu \tag{128}$$

holds for any $\tau_2 \geq \tau_1 \geq 0$.

Proof. Apply the N-estimate for $\Box_g \psi^\tau_\flat = F^\tau_\flat$ in physical space from τ_1 to τ_2. The spacetime term $K^N\left[\psi^\tau_\flat\right]$ has a good sign near the event horizon while away from it it is estimated from Proposition 9.3. Add the estimate (33) and the estimate of Proposition 9.3 to the resulting estimate. This yields (128) except for the errorterm

$$\int_{\mathcal{R}(\tau_1,\tau_2)} F^\tau_\flat N\left(\psi^\tau_\flat\right) \tag{129}$$

on the right hand side. The latter is controlled by borrowing an ϵ from the spacetime term on the left and controlling $\int_{\mathcal{R}(\tau_1,\tau_2)} |F^\tau_\flat|^2$ from the boundedness statement as in Section 9.2. □

Remark 9.5. *Using elementary spectral analysis (involving Weyl's law and compactness arguments) one can show that the first term on the right hand side is always finite (with a constant that may blow up as* $L \to \infty$*). This is not needed in the following, as we are going to apply the above Proposition only for special* Σ_{τ_1}*, for which a better, completely explicit, bound for the right hand side is available.*

9.6 Higher derivatives

The estimates of Propositions 9.3 and 9.4 can be obtained for all higher derivatives in by commuting globally with T and locally near the horizon with the redshift vectorfield (as in [15, 26]) and locally near infinity with angular momentum operators. Since these estimates have become standard after [15] and have appeared in detail previously in the literature, they are omitted.

---

[5] This difficulty is absent in the asymptotically flat case, where one proves a local integrated decay estimate for the full solution. The problem here arises precisely from the fact that we cannot prove an integrated decay estimate for the high-modes!

10 Estimates for $\psi_\sharp^\tau$

Recall the densities (26) and the energies (27).

Proposition 10.1. *We have the estimate*

$$\int_{\mathcal{D}} e_1 \left[\psi_\sharp^\tau\right] \leq \frac{1}{L} E_2 \left[\psi\right] \cdot \tau \tag{130}$$

Proof. From (165) we already have

$$\int_{\mathcal{D}} \sqrt{g} dt^\star dr d\theta d\phi \; e_2 \left[\psi_\sharp^\tau\right] \, . \leq E_2 \left[\psi\right] \cdot \tau \tag{131}$$

Now we can use that $L \leq \lambda_{m\ell} \left(a\omega\right) + a^2\omega^2$ holds for $\psi_\sharp^\tau$. For instance, for $\partial_{\tilde{\phi}}$,

$$\begin{aligned} L \int_{-\infty}^{\infty} dt \int d\sigma_{\mathbb{S}^2} |\partial_{\tilde{\phi}} \psi_\sharp^\tau|_\gamma^2 = L \int_{-\infty}^{\infty} \sum_{m\ell} |im\widehat{\psi_\sharp^\tau}_{m\ell}^{(a\omega)}|^2 d\omega \leq \\ \int_{-\infty}^{\infty} \sum_{m\ell} \left(\lambda_{m\ell} \left(a\omega\right) + a^2\omega^2\right) |\widehat{\left(\partial_{\tilde{\phi}} \psi_\sharp^\tau\right)}_{m\ell}^{(a\omega)} |^2 d\omega \\ \leq C_{M,l,a} \int_{-\infty}^{\infty} dt \int d\sigma_{\mathbb{S}^2} \left[|{}^\gamma\nabla \partial_{\tilde{\phi}} \psi_\sharp^\tau|_\gamma^2 + |\partial_t \partial_{\tilde{\phi}} \psi_\sharp^\tau|^2\right] , \end{aligned} \tag{132}$$

and similarly with the derivatives ∂_r, ∂_θ and ∂_t (and linear combinations like the regular Z-derivative defined in (24)). Multiplying (132) (and its analogues for the other derivatives) with their appropriate r-weight and integrating over the domain of outer communications, we obtain

$$\int_{\mathcal{D}} e_1 \left[\psi_\sharp^\tau\right] \leq \frac{1}{L} \int_{\mathcal{D}} \sqrt{g} dt^\star dr d\theta d\phi \; e_2 \left[\psi_\sharp^\tau\right] \tag{133}$$

from which the claim follows. $\square$

11 Proof of Theorem 1.1: Logarithmic decay

Starting from the estimate

$$\int_{\mathcal{R}(0,\tau/2)} r^2 \sin\theta \; dt^\star dr d\theta d\phi \; e_1 \left[\psi_\flat^\tau\right] \leq \int_{\mathcal{D}} e_1 \left[\psi_\flat^\tau\right] \leq \int_{\mathcal{D}} e_1 \left[\psi^\tau\right] \leq E_1 \left[\psi\right] \cdot \tau \, ,$$

we find a slice $\Sigma_{\tau'}$ with $0 < \tau' < \frac{\tau}{2}$, such that

$$\int_{\Sigma_{\tau'}} r^2 \sin\theta \; dr d\theta d\phi \; e_1 \left[\psi_\flat^\tau\right] \lesssim E_1 \left[\psi\right] . \tag{134}$$

We apply Proposition 9.4 from that slice $\Sigma_{\tau'}$ to any slice to the future of it to obtain in particular

$$\int_{\mathcal{R}(\tau',\tau-1)} r^2 \sin\theta \; dt^\star dr d\theta d\phi \; e_1\left[\psi^\tau_\flat\right] \lesssim C_{M,l,a} \cdot e^{C_{M,l,a} L \frac{\pi}{2}} \int_{t^\star=0} J^N_\mu\left[\psi\right] n^\mu \, .$$

Adding the estimate of Proposition 10.1, we find

$$\int_{\mathcal{R}(\tau/2,\tau-1)} e_1\left[\psi^\tau\right] \leq \int_{\mathcal{R}(\tau/2,\tau-1)} \left(e_1\left[\psi^\tau_\flat\right] + e_1\left[\psi^\tau_\sharp\right]\right) \lesssim \left(e^{C_{M,l,a}L} + \frac{\tau}{L}\right) E_2\left[\psi\right] \, .$$

But the strip we are integrating over has length $\sim \frac{\tau}{2}$ and we can therefore extract a good slice between $\frac{\tau}{2} \leq \tau_g \leq \tau - 1$ where

$$\int_{\Sigma_{\tau_g}} r^2 \sin\theta \; dr d\theta d\phi \; e_1\left[\psi^\tau\right] \lesssim \left(\frac{e^{C_{M,l,a}L}}{\tau} + \frac{1}{L}\right) E_2\left[\psi\right] \, . \tag{135}$$

This holds for any L, in particular for $L = \frac{1}{2C_{M,l,a}} \log(\tau)$ which implies that there is a slice Σ_τ between $\frac{\tau}{2}$ and $\tau - 1$ such that the energy decays like $\frac{1}{\log\tau}$. Hence we have obtained a dyadic sequence of good slices through which the energy decays like $(\log\tau)^{-1}$.

Proposition 11.1. *The non-degenerate energy*

$$f(\tau) = \int_{\Sigma_\tau} e_1\left[\psi\right] \sim \int_{\Sigma_\tau} J^N_\mu\left[\psi\right] n^\mu \tag{136}$$

satisfies

$$f(\tau) \lesssim \frac{1}{\log\tau} \tag{137}$$

for all $\tau \geq 2$.

Proof. Note that the statement is immediate for the degenerate Hawking-Reall energy $\int_{\Sigma_\tau} J^K_\mu n^\mu_\Sigma$ in view of (30): The $(\log\tau)^{-1}$-decay is exported dyadically (from the dyadic sequence of good slices already obtained) implying that the *degenerate* energy decays like $(\log\tau)^{-1}$ through any slice.

This allows us to revisit the estimate for the vectorfield N (cf. 33). Namely, we estimate the error-term in the interior (away from the horizon) using the K-energy which has already been shown to decay logarithmically to obtain the local estimate ($\tau_2 \geq \tau_1 \geq 2$):

$$f(\tau_2) + \int_{\tau_1}^{\tau_2} d\tau f(\tau) \leq C \cdot f(\tau_1) + C \max\left(\left(\tau_2 - \tau_1\right), 1\right) \frac{1}{\log\tau_1} \, . \tag{138}$$

We claim that this estimate implies that

$$f(\tau) \leq \tilde{C} \left(\log\tau\right)^{-1} \tag{139}$$

for $\tau \geq 2$ using a bootstrap argument. Here is a sketch: Wlog, $C \geq 1$. Define $\mathfrak{l} = 8C^2$. Consider the region where $f(\tau) \leq A(\log \tau)^{-1}$ for $A = 32C^3$. Decompose the region into strips of length $\mathfrak{l}$ (the last one may be shorter). In each strip, i.e. between τ_n and τ_{n+1}, we find a good slice for which

$$f(\tau_{good}) \leq \frac{1}{\mathfrak{l}} C \frac{A}{\log \tau_n} + C \frac{1}{\log \tau_n} = \left(\frac{CA}{\mathfrak{l}} + C\right) \frac{1}{\log \tau_n} . \tag{140}$$

Applying the estimate from the good slice up to any slice to the past of the next good slice we find

$$f(\tau) \leq \left(A \frac{C^2}{\mathfrak{l}} + C^2 + C \ 2\mathfrak{l}\right) \frac{1}{\log \tau_n} < \frac{3A}{4 \log \tau_n} < \frac{A}{\log \tau} , \tag{141}$$

where the last step uses that we can assume that $\tau_n \gg \mathfrak{l}$ and hence $\frac{1}{\tau_n} \sim \frac{1}{\tau_{n+2}}$ with a constant very close to 1. This improves the bootstrap assumption. $\square$

12 Some remarks on the Schwarzschild-AdS case

While we are not able to improve the logarithmic decay rate in the case of Schwarzschild-AdS (in accord with the heuristics given in the introduction), one can nevertheless extract some more information about the solution in this special case. The underlying reason is that for $a = 0$ the background is spherically symmetric and one can expand the solution into the familiar spherical harmonics, Y_{lm}, which have eigenvalues $\ell(\ell+1)$ (with a $2\ell + 1$ degeneration for fixed ℓ) *before* cutting off in time. One separates the solution into a part which $\ell(\ell+1) \leq L$ and a part for which $\ell(\ell+1) \geq L$:

$$\psi_\flat\left(t, r, \theta, \tilde{\phi}\right) = \sum_{m\ell} \int_{\mathbb{S}^2} \sin \underline{\theta} d\underline{\theta} d\underline{\tilde{\phi}} \ \psi\left(t, r, \underline{\theta}, \underline{\tilde{\phi}}\right) Y_{\ell m}\left(\underline{\theta}, \underline{\tilde{\phi}}\right) Y_{\ell m}\left(\theta, \tilde{\phi}\right) \chi\left(\frac{\ell(\ell+1)}{L}\right)$$

$$\psi_\sharp\left(t, r, \theta, \tilde{\phi}\right) = \psi\left(t, r, \theta, \tilde{\phi}\right) - \psi_\flat\left(t, r, \theta, \tilde{\phi}\right)$$

where χ is the familiar cut-off functions defined in Section 5.1. Because there is no cut-off in time, both $\psi_\flat$ and $\psi_\sharp$ satisfy the *homogeneous* wave equation. Moreover, since the non-degenerate energies of $\psi_\flat$ are easily seen to be controlled (uniformly in L) by the full ψ-energies, one obtains separate uniform boundedness statements for $\psi_\flat$ and $\psi_\sharp$ for free.[6]

With this in mind, starting from $\Box_g \psi_\flat + \frac{\alpha}{l^2} \psi_\flat = 0$, one cuts $\psi_\flat$ off in time and repeats the previous analysis for $\psi_\flat$. Since the time-cut-off commutes with the cut-off in angular momentum in the spherically-symmetric case, we now obtain the following version of Proposition 9.4:

---

[6] Note that this was not possible in the Kerr-AdS-case where the coupling of the $S_{m\ell}$ with ω prevented one from estimating the $\psi_\flat$-energy on a spacelike slice from the ψ-energy. It is an interesting question whether a more refined spectral analysis of the $S_{m\ell}$ can provide such an estimate.

$$\int_{\Sigma_{\tau_2}} J^N_\mu \left[\psi_\flat\right] n^\mu + \int_{\mathcal{R}(\tau_1,\tau_2)} J^N_\mu \left[\psi_\flat\right] n^\mu \lesssim \int_{\Sigma_{\tau_1}} J^N_\mu \left[\psi_\flat\right] n^\mu$$
$$+ C_{M,l,a} \cdot e^{C_{M,l,a} L \frac{\pi}{2}} \int_{\Sigma_{\tau_1}} J^N_\mu \left[\psi_\flat\right] n^\mu \, . \tag{142}$$

Note that the last term on the right now involves only $\psi_\flat$ and integration over Σ_{τ_1}. The above estimate implies that $\psi_\flat$ decays exponentially:

$$\int_{\Sigma_{t^\star}} J^N_\mu \left[\psi_\flat\right] n^\mu \lesssim \exp\left(-e^{-C_{M,l,a} L} \cdot t^\star\right) \, . \tag{143}$$

Despite this strong decay, interpolating with $\psi_\sharp$ (for which only the boundedness statement is available) one cannot improve the logarithmic decay rate for the full solution. The question whether the low modes in the Kerr-AdS case decay exponentially remains open.

We finally remark that the exponential type multiplier estimates proven are sufficiently robust to establish this type of decay for any stationary spherically-symmetric black hole solution (asymptotically-flat or asymptotically AdS). This is because the exponential estimates only depend on the behavior at the ends, i.e. the asymptotics near the horizon and near infinity. The precise set-up and statements will be elaborated on in future work.

13 Proof of Theorem 1.3: Absence of periodic solutions

We give an elementary proof that there exist no time-periodic solutions for the massive wave equation on Kerr-AdS spacetimes in the parameter range $|a|l < r_+$. For $\alpha = 0$ and the parameters (a, M, l) in the range for which the boundedness theorem holds, this result is an immediate consequence of the $(\log t^\star)^{-1}$-decay established in the main theorem. However, whereas the main theorem required the construction of elaborate multipliers, the exclusion of periodic solutions can be obtained from elementary ODE arguments and the redshift effect near the horizon only. It is for this reason together with the fact that the latter result holds for a wider range of parameters that we include its proof here.

Proof of Theorem 1.3. Step 1: We first prove that, as in the Schwarzschild case (cf. the Appendix of [29]), there exists no *stationary* (with respect to the globally timelike Hawking Reall vectorfield K) solution. This is accomplished by the usual "Bekenstein"-argument [5, 25] combined with a Hardy inequality for the mass term. Assuming that ψ is a solution of $\Box\psi + \frac{\alpha}{l^2}\psi = 0$ which satisfies $K(\psi) = T(\psi) + \lambda\Phi(\psi) = 0$ for $\lambda = \frac{a\Xi}{r_+^2+a^2}$, the wave equation reduces in

Boyer-Lindquist coordinates to:

$$\frac{\Xi}{\Sigma|\sin\theta|}\Big[\partial_r\left(\sqrt{-g}g^{rr}\partial_r\psi\right)+\partial_\theta\left(\sqrt{-g}g^{\theta\theta}\partial_\theta\psi\right)\Big]$$
$$+\left(g^{\tilde{\phi}\tilde{\phi}}-2\lambda g^{t\tilde{\phi}}+\lambda^2 g^{tt}\right)\partial^2_{\tilde{\phi}}\psi+\frac{\alpha}{l^2}\psi=0\,. \tag{144}$$

The expression in the round bracket of the second line is (setting $\det A = g_{tt}g_{\tilde{\phi}\tilde{\phi}}-\left(g_{t\tilde{\phi}}\right)^2=-\frac{\Delta_-\Delta_\theta}{\Xi^2}\leq 0$)

$$P=g^{\tilde{\phi}\tilde{\phi}}-2\lambda g^{t\tilde{\phi}}+\lambda^2 g^{tt}=\frac{1}{detA}\left(g_{tt}+2\lambda g_{t\tilde{\phi}}+\lambda^2 g_{\tilde{\phi}\tilde{\phi}}\right)$$
$$=\frac{1}{detA}g\left(K,K\right)\geq 0\,, \tag{145}$$

since the Hawking-Reall vectorfield is timelike in the black hole exterior. Note that the vector field ∂_r (associated to Boyer-Lindquist coordinates) is given, in terms of $(\partial_{t^\star},\partial_\phi,R)$, where $R(f)=\frac{\partial f}{\partial r}|_{(t^*,\theta,\phi)=const}$ is the radial vector field of the $(t^\star,r,\theta,\phi)$ coordinates, by:

$$\partial_r=\frac{1}{\Delta_-}\left(\frac{2Mr}{1+\frac{r^2}{l^2}}\partial_{t^\star}+a\Xi\partial_\phi\right)+R$$
$$=\frac{1}{\Delta_-}\frac{2Mr}{1+\frac{r^2}{l^2}}K-\frac{2Ma\Xi\left(r-r_+\right)}{\Delta_-\left(\left(r^2+a^2\right)\left(1+\frac{r^2}{l^2}\right)\right)}\partial_\phi+R, \tag{146}$$

where K is the Hawking-Reall vector field. Since $K(\psi)=0$ and since the last two terms are regular vectorfields on the event horizon, it follows that $\partial_r\psi$ extends continously to any sphere arising from the intersection of a $t^*=const$ slice and the future event horizon.

Multiplying (144) by $\frac{\Sigma|\sin\theta|}{\Xi}\psi$, we find after integration by parts on a $t^*=const$ slice[7]:

$$\int_{r=r_+}^{\infty}\int_{\theta,\phi}drd\theta d\phi\Big\{\sqrt{-g}\left(g^{rr}(\partial_r\psi)^2+g^{\theta\theta}(\partial_\theta\psi)^2+P(\partial_{\tilde{\phi}}\psi)^2\right)$$
$$-\frac{\alpha}{l^2}\psi^2\frac{\Sigma|\sin\theta|}{\Xi}\Big\}(t^*,r,\theta,\phi)=0, \tag{147}$$

where the boundary term at infinity has vanished in view of the decay of ψ and the boundary term at the horizon vanished since g^{rr} extends continuously to 0 at the event horizon and since $\psi\partial_r\psi$ is regular at r_+, as established above.

As in [26, 29], using the bound $|a|<\frac{l}{2}$, the zero order term can be absorbed into the first term via a Hardy inequality, from which it follows that $\psi=0$.

---

[7] The integration is performed on $t^*=const$ slice to avoid the bifurcate sphere $\{t=const\}\cap\{r=r_+\}$ where all the coordinate systems that we have so far introduced break down.

Step 2: We use a microlocal decomposition to reduce the analysis to that of an ode with trivial data. More precisely, in view of the separability of the geodesic equation, we may restrict our attention to solutions ψ of the form:

$$\psi = e^{-i\omega t} f(r^*) S_{m\ell}(a\omega, \cos\theta) e^{im\tilde{\phi}}, \tag{148}$$

where $S_{m\ell}(a\omega, \cos\theta)e^{im\tilde{\phi}}$ are the (modified) oblate spheroidal harmonics introduced in Section 5.2. In view of Step 1 we can assume $\omega - \omega_+ \neq 0$. Since ψ is regular at the future event horizon and since the vector field $-\Delta_- Z = \partial_t + \frac{a\Xi}{r_+^2+a^2}\partial_\phi - \partial_{r^*}$ vanishes on $\mathcal{H}^+$ (cf. (24), it follows that:

$$f' + i\left(\omega - \frac{a\Xi m}{r_+^2 + a^2}\right) f = 0.$$

In fact, we have on any $t^* = const$ slice, $-\Delta_- Z = \Delta_- D$ where D is a regular vector field at the horizon. Hence we obtain that

$$f' + i\left(\omega - \frac{a\Xi m}{r_+^2 + a^2}\right) f = \mathcal{O}\left(\Delta_-\right),$$

near the horizon. Since $\frac{dr}{dr^*} = \frac{\Delta_-}{r^2+a^2}$ vanishes at the horizon, the rescaled function $u = \sqrt{r^2+a^2} f$ also satisfies:

$$u' + i\left(\omega - \frac{a\Xi m}{r_+^2 + a^2}\right) u = \mathcal{O}\left(\Delta_-\right). \tag{149}$$

Moreover, since ψ is in the class CH^1_{AdS}, we have the following uniform[8] radial decay estimates (recall that $'$ denotes r^*-differentiation):

$$\psi r^{3/2} = o(1) \quad , \quad \psi' r^{1/2} = o(1). \tag{150}$$

Hence, we find for $u = \sqrt{r^2+a^2}\psi$, the decay rates

$$u r^{1/2} = o(1) \quad , \quad u' r^{-1/2} = o(1). \tag{151}$$

In Section 5.3, we obtained the following ode (52) for each $u^{(a\omega)}_{m\ell}$ (which we will write short hand as u)

$$u'' + \left(\omega^2 - V\left(r\right)\right) u = H. \tag{152}$$

Compared to (52), here we have $H = 0$ since no cut-off is needed in view of the ansatz (148). Recall the current (76). From the decay of u at ∞, it follows that

$$Q_T[u] = 0 \qquad \text{and hence} \quad Im\left(u'\bar{u}\right) = 0 \tag{153}$$

---

[8] A priori, for a general solution, the r decay is only locally uniform in the t variable. Here, of course, we have global uniform radial decay in view of the ansatz.

globally, which in conjunction with the boundary condition (149) on the horizon implies that u vanishes on the horizon.

Next we choose the free function z in the red-shift current (77) to be $z = V_{red}^{-1}$, where V_{red} was defined in (64). Since

$$\left|u' + i\left(\omega - \omega_+\right) u\right|^2 = \mathcal{O}(r - r_+)^2$$

from (149) and $u = 0$ on the horizon, it follows that $Q_{red}^{V_{red}^{-1}} = 0$ on the horizon. The derivative satisfies (use (153))

$$Q'_{red} = -\frac{V'_{red}}{V_{red}^2}\left|u' + i\left(\omega - \omega_+\right) u\right|^2 - \frac{1}{V_{red}}(|u|^2)'\left(V_{red} - \tilde{V}\right) \leq 0 \qquad (154)$$

in a neighborhood of the horizon, which implies that $Q_{red}^{V_{red}^{-1}}$ is non-increasing near the horizon. On the other hand, one easily checks that $Q_{red}^{V_{red}^{-1}} \geq 0$ holds near the horizon, which is only consistent with (154) if $Q_{red}^{V_{red}^{-1}}$ and hence u are identically zero in a neighbourhood of the horizon. Solving the linear ode (152) with trivial initial conditions we conclude $u = 0$ identically. □

14 Acknowledgments

The authors thank Mihalis Dafermos and Igor Rodnianski for insightful discussions. We also acknowledge the hospitality of the Erwin Schrödinger Institute, Vienna, where part of this work was being completed in July 2011 during the workshop "Dynamics of General Relativity". J.S. thanks the Albert Einstein Institute, Potsdam for financial support during his postdoctoral fellowship.

A Basic Fourier estimates

Let $\varphi\left(t, r, \theta, \tilde{\phi}\right)$ be a smooth function in Boyer-Lindquist coordinates and assume that φ is also L^2 in time. We have the Fourier-transformation of φ,

$$\widehat{\varphi}\left(\omega, r, \theta, \tilde{\phi}\right) = \frac{1}{\sqrt{2\pi}}\int_{-\infty}^{\infty} dt\ e^{i\omega t}\varphi\left(t, r, \theta, \tilde{\phi}\right) . \qquad (155)$$

We can, for each ω and r, interpret $\widehat{\varphi}$ as a function on the Boyer-Lindquist spheres $\mathbb{S}^2$. There is an orthonormal basis of L^2-eigenfunctions $S_{m\ell}\left(\theta, \tilde{\phi}\right) = S_{m\ell}\left(a\omega, \cos\theta\right) e^{im\tilde{\phi}}$ on these spheres, which is obtained as the solutions of (42). Any (say smooth) function on the sphere can be expanded in terms of these eigenfunctions. Completeness and orthogonality of this basis yield:

$$\widehat{\varphi}\left(\omega, r, \theta, \tilde{\phi}\right) = \sum_{ml} \widehat{\varphi}_{m\ell}^{(a\omega)}\left(r\right) S_{m\ell}\left(\theta, \tilde{\phi}\right) , \qquad (156)$$

$$\widehat{\varphi}^{(a\omega)}_{m\ell}(r) = \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \; \widehat{\varphi}\left(\omega, r, \theta, \tilde{\phi}\right) S^{\star}_{m\ell}\left(\theta, \tilde{\phi}\right) \,, \tag{157}$$

where $\sqrt{\gamma} = \sin\theta$ denotes the volume element with respect to which the orthogonality relation of the $S_{m\ell}$ holds, cf. again (42). We have the important Parseval identity on $\mathbb{S}^2$,

$$\int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \; |\widehat{\varphi}\left(\omega, r, \theta, \tilde{\phi}\right)|^2 = \sum_{m\ell} |\widehat{\varphi}^{(a\omega)}_{m\ell}(r)|^2 \,. \tag{158}$$

Note that both sides depend on ω.

In the paper, we restrict to low modes. More precisely, we define for given L, the quantity

$$\widehat{\varphi}^{(a\omega)}_{m\ell}\Big|_{\flat}(r) = \widehat{\varphi}^{(a\omega)}_{m\ell}(r) \cdot \zeta\left(\frac{a^2\omega^2 + \lambda_{m\ell}}{L}\right) \tag{159}$$

and the associated quantity

$$\widehat{\varphi}_{\flat}\left(\omega, r, \theta, \tilde{\phi}\right) = \sum_{ml} \widehat{\varphi}^{(a\omega)}_{m\ell}|_{\flat}(r) \, S_{m\ell}\left(\theta, \tilde{\phi}\right) \,. \tag{160}$$

Then, again by Parseval and the fact that $|\zeta| \leq 1$, we have

$$\begin{aligned} \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \; |\widehat{\varphi}_{\flat}\left(\omega, r, \theta, \tilde{\phi}\right)|^2 &= \sum_{m\ell} |\widehat{\varphi}^{(a\omega)}_{m\ell}|_{\flat}(r)|^2 \\ \leq \sum_{m\ell} |\widehat{\varphi}^{(a\omega)}_{m\ell}(r)|^2 &= \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \; |\widehat{\varphi}\left(\omega, r, \theta, \tilde{\phi}\right)|^2 \end{aligned}$$

Integrating this identity in ω and using that $\sqrt{\gamma}$ is independent of ω, we obtain (using Parseval in t and ω) that

$$\begin{aligned} &\int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \; |\varphi_{\flat}\left(t, r, \theta, \tilde{\phi}\right)|^2 \\ \leq &\int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \; |\varphi\left(t, r, \theta, \tilde{\phi}\right)|^2 \,. \end{aligned} \tag{161}$$

Of course, the same identity holds for $\varphi_{\sharp}$.

Applications

- Letting $\varphi = \psi^{\tau}$ we can apply the estimate (161) for fixed $r > r_+$ corresponding to $R^{\star}_{-\infty}$. In the limit $R^{\star}_{-\infty} \to -\infty$ one obtains an integral over the event horizon:

$$\begin{aligned} &\lim_{R^{\star}_{-\infty} \to -\infty} \int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \; |\psi^{\tau}_{\flat}\left(t, r\left(R^{\star}_{-\infty}\right), \theta, \tilde{\phi}\right)|^2 \\ \leq &\lim_{R^{\star}_{-\infty} \to -\infty} \int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \; |\psi^{\tau}\left(t, r\left(R^{\star}_{-\infty}\right), \theta, \tilde{\phi}\right)|^2 \\ &\leq \int_{\mathcal{H}(0,\tau)} \sqrt{\gamma} dt^{\star} d\theta d\tilde{\phi} \; |\psi^{\tau}\left(t, r_+, \theta, \tilde{\phi}\right)|^2 \,. \end{aligned} \tag{162}$$

- Letting $\varphi = F$, the inhomogeneity in the wave equation for $\psi^\tau_\flat$, which is compactly supported in time (in two strips, actually). Integrating (161) over $\int_{R^\star_{-\infty}}^{R^\star_\infty} dr^\star$, we obtain that

$$\int_{\mathcal{R}} r^2 |F_\flat|^2 \leq \int_{\mathcal{R}} r^2 |F|^2 = \int_{\mathcal{R}(0,1)\cup\mathcal{R}(\tau,\tau+1)} r^2|F|^2 \lesssim \|\psi\|^2_{H^1_{AdS}} \, . \tag{163}$$

- The estimate (161) also holds for derivatives. For instance,

$$\begin{aligned}
&\int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \, \left| \left(-\partial_t + \partial_{r^\star} - \frac{a\Xi}{r_+^2+a^2}\partial_{\tilde{\phi}}\right) \varphi_\flat \right|^2 \\
&= \int_{-\infty}^{\infty} d\omega \sum_{m\ell} \left| \left[\left(-\partial_t + \partial_{r^\star} - \frac{a\Xi}{r_+^2+a^2}\partial_{\tilde{\phi}}\right) \varphi_\flat \right]^{a\omega}_{m\ell} \right|^2 \\
&= \int_{-\infty}^{\infty} d\omega \sum_{m\ell} \left| i\omega \widehat{\varphi}^{(a\omega)}_{m\ell}|_\flat + \partial_{r^\star}\widehat{\varphi}^{(a\omega)}_{m\ell}|_\flat - i\frac{a\Xi m}{r_+^2+a^2}\widehat{\varphi}^{(a\omega)}_{m\ell}|_\flat \right|^2 \\
&= \int_{-\infty}^{\infty} d\omega \sum_{m\ell} \left[\zeta\left(\frac{a^2\omega^2+\lambda_{m\ell}}{L}\right)\right]^2 \left| i\omega \widehat{\varphi}^{(a\omega)}_{m\ell} + \partial_{r^\star}\widehat{\varphi}^{(a\omega)}_{m\ell} - i\frac{a\Xi m}{r_+^2+a^2}\widehat{\varphi}^{(a\omega)}_{m\ell} \right|^2 \\
&\leq \int_{-\infty}^{\infty} d\omega \sum_{m\ell} \left| i\omega \widehat{\varphi}^{(a\omega)}_{m\ell} + \partial_{r^\star}\widehat{\varphi}^{(a\omega)}_{m\ell} - i\frac{a\Xi m}{r_+^2+a^2}\widehat{\varphi}^{(a\omega)}_{m\ell} \right|^2 \\
&= \int_{-\infty}^{\infty} d\omega \sum_{m\ell} \left| \left[\left(-\partial_t + \partial_{r^\star} - \frac{a\Xi}{r_+^2+a^2}\partial_{\tilde{\phi}}\right) \varphi \right]^{a\omega}_{m\ell} \right|^2 \\
&= \int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \, \left| \left(-\partial_t + \partial_{r^\star} - \frac{a\Xi}{r_+^2+a^2}\partial_{\tilde{\phi}}\right) \varphi \right|^2 .
\end{aligned}$$

Higher derivatives can be dealt with similarly. We summarize this as

Lemma A.1. *Let D denote any operator of the collection ∂_t, $\partial_{\tilde{\phi}}$, ∂_θ, $\partial_{r^\star}$ and Z (defined in (24)) of Boyer-Lindquist derivative operators. For a non-negative integer k let D^k denote any k^{th}-order combination of these derivatives. Then, for any k,*

$$\int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \, |D^k \varphi_\flat|^2 \leq \int_{-\infty}^{\infty} dt \int_{\mathbb{S}^2} \sqrt{\gamma} d\theta d\tilde{\phi} \, |D^k \varphi|^2 \, . \tag{164}$$

Corollary A.2. *Letting $\varphi = \psi^\tau$, we have for any weight function $w(r)$*

$$\int_{\mathcal{D}} w(r) \, \sin\theta \, dt dr^\star d\theta d\tilde{\phi} \, \left| D^k \psi^\tau_\flat \right|^2 \leq \int_{\mathcal{D}} w(r) \, \sin\theta \, dt dr^\star d\theta d\tilde{\phi} \, \left| D^k \psi^\tau \right|^2 .$$

We may now choose $w(r)$ with the correct weights near the horizon and near infinity to mimic the volume element $\sqrt{g}$. Since the derivatives D are all regular, and can readily be converted into the regular $(t^\star, r, \theta, \phi)$- coordinates, we finally obtain (using that the right hand side is only supported for $0 \leq t^\star \leq \tau$):

$$\int_{\mathcal{D}} \sqrt{g} dt^\star dr d\theta d\phi \; e_2\left[\psi^\tau_\flat\right] \leq \int_{\mathcal{R}(0,\tau)} \sqrt{g} dt^\star dr d\theta d\phi \; e_2\left[\psi^\tau\right] \leq E_2\left[\psi\right] \tau \quad (165)$$

The same estimate holds for $\psi^\tau_\sharp$.

References

[1] Anderson, M. T. On the uniqueness and global dynamics of AdS spacetimes. *Class. Quant. Grav. 23* (2006), 6935–6954, hep-th/0605293.

[2] Andersson, L., and Blue, P. Hidden symmetries and decay for the wave equation on the Kerr spacetime. arXiv:0908.2265.

[3] Aretakis, S. Decay of Axisymmetric Solutions of the Wave Equation on Extreme Kerr Backgrounds. arXiv:1110.2006.

[4] Bachelot, A. The Dirac System on the Anti-de Sitter Universe. *Commun. Math. Phys. 283* (2008), 127–167.

[5] Bekenstein, J. D. Nonexistence of baryon number for static black holes. *Phys.Rev. D5* (1972), 1239–1246.

[6] Bizon, P., and Rostworowski, A. On weakly turbulent instability of anti-de Sitter space. *Phys.Rev.Lett. 107* (2011), 031102, 1104.3702.

[7] Bony, J.-F. and Häfner, D. Decay and non-decay of the local energy for the wave equation in the De Sitter-Schwarzschild metric. *Comm. Math. Phys. 282* (2008), 697–719.

[8] Breitenlohner, P., and Freedman, D. Z. Stability in Gauged Extended Supergravity. *Ann. Phys. 144* (1982), 249.

[9] Burq, N. Décroissance de l'énergie locale de l'équation des ondes pour le problème extérieur et absence de résonance au voisinage du réel. *Acta Math. 180*, 1 (1998), 1–29.

[10] Cardoso, V., and Dias, O. J. Small Kerr-anti-de Sitter black holes are unstable. *Phys.Rev. D70* (2004), 084011, hep-th/0405006.

[11] Cardoso, V., Dias, O. J., Lemos, J. P., and Yoshida, S. The Black hole bomb and superradiant instabilities. *Phys.Rev. D70* (2004), 044039, hep-th/0404096.

[12] Cardoso, V., Dias, O. J., and Yoshida, S. Classical instability of Kerr-AdS black holes and the issue of final state. *Phys.Rev. D74* (2006), 044008, hep-th/0607162.

[13] Dafermos, M. *The Black Hole Stability problem.* Newton Institute, Cambridge, http://www.newton.ac.uk/webseminars/pg+ws/2006/gmx/1010/dafermos/, 2006.

[14] Dafermos, M., and Rodnianski, I. A proof of Price's law for the collapse of a self- gravitating scalar field. *Invent. Math. 162* (2005), 381–457, gr-qc/0309115.

[15] Dafermos, M., and Rodnianski, I. Lectures on black holes and linear waves. *Institut Mittag-Leffler Report no. 14, 2008/2009* (2008), arXiv:0811.0354.

[16] Dafermos, M., and Rodnianski, I. A new physical-space approach to decay for the wave equation with applications to black hole spacetimes. *XVI International Congress on Mathematical Physics, P. Exner (ed), World Scientific, London* (2009), arXiv:0910.4957v1.

[17] Dafermos, M., and Rodnianski, I. The red-shift effect and radiation decay on black hole spacetimes. *Comm. Pure Appl. Math. 62* (2009), 859–919, gr-qc/0512119.

[18] Dafermos, M., and Rodnianski, I. Decay for solutions of the wave equation on Kerr exterior spacetimes I-II: The cases $|a| \ll M$ or axisymmetry. arXiv:1010.5132.

[19] Dafermos, M., and Rodnianski, I. The black hole stability problem for linear scalar perturbations. *to appear in Proceedings of the* 12^{th} *Marcel Grossmann Meeting* (2010), arXiv:1010.5137.

[20] Dafermos, M., and Rodnianski, I. A proof of the uniform boundedness of solutions to the wave equation on slowly rotating Kerr backgrounds. *Invent. Math. 185 (3)* (2011), 467–559, arXiv:0805.4309.

[21] Dyatlov, S. Asymptotic distribution of quasi-normal modes for Kerr-de Sitter black holes. arXiv:1101.1260.

[22] Dyatlov, S. Quasi-normal modes and exponential energy decay for the Kerr-de Sitter black hole. *Comm. Math. Phys. 306* (2011), 119–163, arXiv:1003.6128.

[23] Festuccia, G., and Liu, H. A Bohr-Sommerfeld quantization formula for quasinormal frequencies of AdS black holes. *Advanced Science Letters 2 (2)* (2009), 221–235, arXiv:0811.1033.

[24] Hawking, S. W., and Reall, H. S. Charged and rotating AdS black holes and their CFT duals. *Phys. Rev. D61* (2000), 024014, hep-th/9908109.

[25] Heusler, M. *Black Hole Uniqueness Theorems.* Cambridge Lecture Notes in Physics, Cambridge, 1996.

[26] Holzegel, G. On the massive wave equation on slowly rotating Kerr-AdS spacetimes. *Comm. Math. Phys. 294* (2010), 169–197, arXiv:0902.0973.

[27] Holzegel, G. Well-posedness for the massive wave equation on asymptotically anti-de-Sitter spacetimes. *to appear in Journal of Hyperbolic Differential Equations* (2011), arXiv:1103.0710.

[28] Holzegel, G., and Smulevici, J. Self-gravitating Klein-Gordon fields in asymptotically Anti-de Sitter spacetimes. *to appear in Annales Henri Poincaré* (2011), arXiv:1103.0712.

[29] Holzegel, G., and Smulevici, J. Stability of Schwarzschild-AdS for the spherically symmetric Einstein-Klein Gordon system. arXiv:1103.3672.

[30] Horowitz, G. T., and Hubeny, V. E. Quasinormal modes of AdS black holes and the approach to thermal equilibrium. *Phys.Rev. D62* (2000), 024027, hep-th/9909056.

[31] J. Marzuola, J. Metcalfe, D. Tataru, M. Tohaneanu. Strichartz Estimates on Schwarzschild Black Hole Backgrounds. *Comm. Math. Phys. 293* (2010), arXiv:0802.3942.

[32] R. Melrose, A. Sá Barreto and A. Vasy. Asymptotics of solutions of the wave equation on de Sitter-Schwarzschild space. arXiv:0811.2229.

[33] Tataru, D., and Tohaneanu, M. Local energy estimate on Kerr black hole backgrounds. *Int. Math. Res. Not. 2011* (2011), 248–292, arXiv:0810.5766.

[34] Vasy, A. The wave equation on asymptotically Anti-de Sitter spaces. *to appear in Analysis and PDE* (2009), arXiv:0911.5440.

[35] Vasy, A. The wave equation on asymptotically de Sitter-like spaces. *Advances in Mathematics 223* (2010), 49–97, arXiv:0706.3669.

[36] Vasy, A. and Dyatlov, S. Microlocal analysis of asymptotically hyperbolic and Kerr-de Sitter spaces. arXiv:1012.4391.